

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

12

EUROPEAN PATENT APPLICATION

21 Application number: 83305650.0

22 Date of filing: 22.09.83

61 Int. Cl.³: **C 07 C 69/738**

C 07 C 59/82, C 07 C 177/00
C 07 C 67/327, C 07 C 51/00
A 61 K 31/557, A 61 K 31/19
A 61 K 31/215

30 Priority: 07.10.82 JP 175267/82
07.10.82 JP 175268/82
14.02.83 JP 21617/83
10.03.83 JP 38190/83

43 Date of publication of application:
25.04.84 Bulletin 84/17

84 Designated Contracting States:
CH DE FR GB IT LI

71 Applicant: **TEIJIN LIMITED**
11, 1-Chome, Minamihonmachi Higashi-ku
Osaka-shi Osaka(JP)

72 Inventor: **Noyori, Ryoji**
136-417, Umemori Shinden
Nisshin-cho Aichi-gun Aichi-ken(JP)

72 Inventor: **Fukushima, Masanori**
1-22, Shinpo-cho
Chikusa-ku Nagoya-shi Aichi-ken(JP)

72 Inventor: **Kurozumi, Seizi**
1-2-28, Higashitokura
Kokubunji-shi Tokyo(JP)

72 Inventor: **Sugura, Satoshi**
Teijin Musashinoryo, 3-5-18
Tamadaira Hino-shi Tokyo(JP)

74 Representative: **Myerscough, Philip Boyd et al,**
J.A.Kemp & Co. 14, South Square Gray's Inn
London, WC1R 5EU(GB)

54 Novel 5-membered cyclic compounds, process for production thereof, and pharmaceutical use thereof.

57 Novel 5-alkylidene-4-substituted-2-cyclopentenones and 5-(1-hydroxy-aliphatic hydrocarbon)-4-substituted-2-cyclopentenones.

These novel cyclopentenones and 5-alkylidene-3-hydroxy-4-substituted cyclopentanones have a pharmaceutical activity for treatment of tumors.

EP 0 106 576 A2

- 1 -

TITLE: "NOVEL 5-MEMBERED CYCLIC COMPOUNDS, PROCESS FOR PRODUCTION THEREOF, AND PHARMACEUTICAL USE THEREOF"

This invention relates to novel 5-membered cyclic compounds, a process for production thereof, and a pharmaceutical use thereof.

Prostaglandin (PG) is a living body regulating substance which is involved in various biological reactions such as contraction of smooth muscles, lowering of blood pressures and inhibition of platelet aggregation. It has been anticipated that prostaglandin as a living body regulating substance affects proliferation of cells. Santoro et al. reported in Cancer. Res. 37, 3774 (1977) that PGE series inhibit growth of tumors of B16 melanoma cell in vivo. W. A. Turner et al. reported that PGA series inhibit tumor cell proliferation and induce differentiation as a result of experiments in vitro using B16 melanoma cell and mouse neuroblastoma [Prostaglandins and Cancer: First International Conference, pages 365-368 (1982)].

Honn et al. reported a possibility of using PGA series as antitumor agents in view of the fact that PGA series strongly inhibit synthesis of DNA [Biochem. Biophys. Res. Commun. 87, 795 (1979)].

M. Fukushima et al. examined the effect of PGD₂ to inhibit proliferation of mouse leukemia cell L1210 and human leukemia cell lines, and reported that IC₅₀ of PGD₂ on L1210 cell is 2.4 micrograms/ml [Biochem. Biophys. Res. Commun., 105, 956 (1982)].

It is an object of this invention to provide novel 5-membered cyclic compounds.

Another object of this invention is to provide novel 5-membered cyclic compounds which have a 5-membered cyclic ring like certain kinds of prostaglandin.

Still another object of this invention is to provide a pharmaceutical use of the 5-membered cyclic

- 2 -

compounds of the invention, especially their use as an antitumor agent.

Still another object of this invention is to provide novel 5-membered cyclic compounds which have better antitumor activity than hitherto known PG series.


Still another object of this invention is to provide novel 5-membered cyclic compounds as highly safe substances with antitumor activity which have no effect on normal cells and do not exhibit any significant toxicity.

Still another object of this invention is to provide novel 5-membered cyclic compounds which show nearly selective antitumor activity as a biological activity and do not substantially show hypotensive activity or platelet aggregation inhibiting activity exhibited by known PGA_2 , and which are therefore very suitable for use as an antitumor agent.

Still another object of this invention is to provide a very simple process for producing the 5-membered cyclic compounds of the invention.

Further objects and advantages of this invention will become apparent from the following description.

As the novel cyclic 5-membered compounds of the invention, the present invention provides a 4,5-disubstituted-2-cyclopentenone selected from the group consisting of 5-alkylidene-4-substituted-2-cyclopentenones represented by the following formula (I)



... (I)

wherein W represents an aliphatic hydrocarbon

- 3 -

group having 1 to 12 carbon atoms which may have a substituent, and Y represents an aliphatic hydrocarbon group having 1 to 12 carbon atoms which may have a substituent,


- 5 and 5-(1-hydroxy aliphatic hydrocarbon)-4-substituted-2-cyclopentenones represented by the following formula (II)



- 10 wherein W' and Y' are the same as W and Y above respectively.

According to this invention, the 5-alkylidene-4-substituted-2-cyclopentenones of formula (I) can be produced by subjecting 5-alkylidene-3-hydroxy-4-substituted-cyclopentanones represented by the following formula (III)

15




- wherein W' and Y' are the same as W and Y above respectively,
- 20 to dehydration reaction, and as required, subjecting the resulting product to a deprotecting, hydrolyzing or salt-forming reaction.

Some of the compounds of formula (III) are disclosed in European Laid-Open Patent Publication No. 0079733 and U. S. Patent Application Serial No. 25 438,379, and are known.


The compounds of formula (III) can be produced

- 4 -

by converting a 3-hydroxy-5-(1-hydroxy-aliphatic hydrocarbon)-4-substituted-cyclopentanone represented by the following formula (IV)



- 5 wherein W'' and Y'' are the same as W and Y
above, respectively,
to a corresponding 3-(t-butyldimethylsilyloxy)-5-
(1-methanesulfonyloxy-aliphatic hydrocarbon)-4-
substituted cyclopentanone, thereafter removing the
10 methanesulfonyloxy group as methanesulfonic acid from
the resulting cyclopentanone and further removing the
t-butyldimethylsilyl group from it. Some of the
compounds of formula (IV) are disclosed in European
Laid-Open Patent Publications Nos. 0019475 and
15 0079733, U. S. Patent No. 4,315,032 and U. S. Patent
Application Serial No. 438379. The above process for
producing the compound of formula (III) from the
compound of formula (IV) is substantially the same as
the methods described in European Laid-Open Patent
20 Publication No. 0079733 and U.S. Patent Application
Serial No. 438,378. Accordingly, the disclosures of
the above-cited European Laid-Open Patent Publications
and U. S. Patent and U. S. Patent Application are
incorporated herein as reference.
- 25 According to the present invention, the 5-
(1-hydroxy-aliphatic hydrocarbon)-4-substituted-2-
cyclopentenones of formula (II) can be produced by
subjecting a 5-(1-hydroxy-aliphatic hydrocarbon)-
3-hydroxy-4-substituted-cyclopentanone represented by
30 the following formula (IV)



... (IV)

wherein W'' and Y'' are the same as W and Y above respectively,

to dehydration reaction, and as required, subjecting the resulting product to a deprotecting, hydrolyzing or salt-forming reaction.

Now, the first process of this invention starting from the 5-alkylidene-3-hydroxy-4-substituted-cyclopentanone of formula (III) and the second process of the invention starting from the 5-(alpha-hydroxy-aliphatic hydrocarbon)-3-hydroxy-4-substituted cyclopentanone of formula (IV) will be described in detail.

In the first process, W' in formula (III) defining the starting material is an aliphatic hydrocarbon group having 1 to 12 carbon atoms, and Y' is likewise an aliphatic hydrocarbon group having 1 to 12 carbon atoms. These aliphatic hydrocarbon groups may be substituted.

The aliphatic hydrocarbon groups for W' and Y' may be linear, branched or cyclic or may contain a carbon-carbon double or triple bond.

Preferably, the aliphatic hydrocarbon groups include, for example, linear or branched C₁₋₁₂ alkyl, alkenyl or alkynyl groups, and cycloalkyl groups having 3 to 8 carbon atoms.

Specific examples of alkyl groups having 1 to 12 carbon atoms are methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, n-decyl, n-undecyl and n-dodecyl.

- 6 -

Specific examples of the alkenyl groups having 1 to 12 carbon atoms are ethenyl, 1-propen-1-yl, 2-propen-1-yl, 1-buten-1-yl, 1,3-butadien-1-yl, 2-buten-1-yl, 1-penten-1-yl, 2-penten-1-yl, 1-hexen-1-yl, 2-hexen-1-yl, 1,5-hexadien-1-yl, 3-hexen-1-yl, 1-hepten-1-yl, 1-octen-1-yl, 1,7-octadien-1-yl, 1-nonen-1-yl, 1-decen-1-yl, 1-undecen-1-yl and 1-dodecen-1-yl.

Specific examples of the alkynyl groups having 1 to 12 carbon atoms are ethynyl, 1-propyn-1-yl, 2-propyn-1-yl, 1-butyn-1-yl, 3-buten-1-yne-1-yl, 2-butyn-1-yl, 1-pentyn-1-yl, 2-pentyn-1-yl, 1-hexyn-1-yl, 2-hexyn-1-yl, 5-hexen-1-yne-1-yl, 3-hexyn-1-yl, 1-heptyn-1-yl, 1-octyn-1-yl, 7-octen-1-yne-1-yl, 1-nonyn-1-yl, 1-decyn-1-yl, 1-undecyn-1-yl and 1-dodecyn-1-yl.

Examples of the cycloalkyl groups having 3 to 8 carbon atoms are cyclopropyl, cyclopentyl, cyclohexyl, cyclooctyl and cyclohexenyl.

These aliphatic hydrocarbon groups may have substituents.

Examples of the substituents include groups of the formula $-\text{COOR}^4$ (wherein R^4 represents a hydrogen atom, an alkyl group having 1 to 10 carbon atoms or one equivalent of a cation); groups of the formula $-\text{OR}^5$ (wherein R^5 represents a hydrogen atom, a C_{1-6} alkyl which may be substituted by a halogen atom, a C_{1-7} carboacyl group, or a phenyl group which may be substituted by a halogen atom, an alkyl group having 1 to 4 carbon atoms or an alkoxy group having 1 to 4 carbon atoms); a phenyl group which may be substituted by a halogen atom, an alkyl group having 1 to 4 carbon atoms, or an alkoxy group having 1 to 4 carbon atoms; and cycloalkyl groups having 3 to 8 carbon atoms which may be substituted by a halogen atom, an alkyl group having 1 to 4 carbon

- 7 -

atoms or an alkoxy group having 1 to 4 carbon atoms.

Specific examples of the groups of the formula $-\text{COOR}^4$ are those in which R^4 is the same alkyl

- group as above having 1 to 10 carbon atoms, or one
5 equivalent of a cation, for example an ammonium
cation such as NH_4^+ , tetramethyl ammonium, mono-
methyl ammonium, dimethyl ammonium, trimethyl ammonium,
benzyl ammonium, phenethyl ammonium, morpholinium
cation, monoethanol ammonium or piperidinium cation,
10 an alkali metal cation such as Na^+ or K^+ , or a
divalent or trivalent metal cation such as $\frac{1}{2}\text{Ca}^{2+}$,
 $\frac{1}{2}\text{Mg}^{2+}$, $\frac{1}{2}\text{Zn}^{2+}$ or $\frac{1}{3}\text{Al}^{3+}$.

- Specific examples of the groups of the formula $-\text{OR}^5$ include a hydroxyl group; alkoxy groups having
15 1 to 6 carbon atoms such as methoxy, ethoxy, n-
propoxy, isopropoxy, n-butoxy, n-pentoxy and n-
hexoxy; carboacyloxy groups having 1 to 7 carbon
atoms such as acetoxo, propionyloxy, n-butyryloxy,
isobutyryloxy, n-valeryloxy, isovaleryloxy, caproyl-
20 oxy, enanthyloxy and benzoyloxy; and a phenoxy group.
The C_{1-6} alkoxy groups for $-\text{OR}^5$ may be substi-
tuted by halogen atoms, thus providing chloromethoxy,
dichloromethoxy, trifluoromethoxy, etc. The phenyl
moiety of the phenoxy group for $-\text{OR}^5$ may be sub-
25 stituted by a halogen atom such as chloro, bromo or
fluoro, an alkyl group having 1 to 4 carbon atoms
such as methyl, ethyl, propyl or butyl, or an alkoxy
group having 1 to 4 carbon atoms such as methoxy,
ethoxy, propoxy or butoxy.


- 30 A phenyl group or a cycloalkyl group having 3
to 8 carbon atoms may also be substituents for the
aforesaid aliphatic hydrocarbon groups. The phenyl
group and the C_{3-8} cycloalkyl group may be substi-
tuted by the same substituents as described above,
35 i.e. a halogen atom, an alkyl group having 1 to 4
carbon atoms or an alkoxy group having 1 to 4 carbon

- 8 -

atoms.


According to this invention, the aforesaid first process which comprises subjecting the 5-alkylidene-3-hydroxy-4-substituted cyclopentenone of formula (III) to dehydration reaction and as required subjecting the resulting product to a deprotecting, hydrolyzing or salt-forming reaction is preferably performed by producing a 5-alkylidene-4-substituted-2-cyclopentenone represented by the following formula

10 (I)-1



... (I)-1

wherein R^1 represents a hydrogen atom or an aliphatic hydrocarbon group having 1 to 10 carbon atoms which may have a substituent, R^2 represents a hydrogen atom, or an aliphatic hydrocarbon group having 1 to 9 carbon atoms which may have a substituent, R^3 represents a hydrogen atom, a hydroxyl group, or a protected hydroxyl group, and the symbol \equiv represents a single, double or triple bond, from a 5-alkylidene-3-hydroxy-4-substituted cyclopentenone represented by the following formula (III)-1



... (III)-1

wherein R^1 , R^2 , R^3 , and the symbol \equiv are the same as given for formula (I)-1.

25

The dehydration reaction of the 5-alkylidene-

- 9 -


3-hydroxy-4-substituted cyclopentenone of formula (III) or (III)-1 is preferably carried out in the presence of a dehydrating agent. Examples of the dehydrating agent include inorganic acids such as hydrochloric acid, hydrobromic acid, hydrofluoric acid and phosphoric acid, organic carboxylic acids such as propionic acid, oxalic acid, citric acid and maleic acid, and organic sulfonic acids such as methanesulfonic acid, ethanesulfonic acid, benzene-sulfonic acid and p-toluenesulfonic acid. Of these, the inorganic acids and organic carboxylic acids are preferred. The amount of the dehydrating agent used is preferably 0.5 to 100 moles, especially preferably 1 to 50 moles, per mole of the 5-alkylidene-3-hydroxy-4-substituted cyclopentanone. As a reaction solvent, there may be used an ether such as tetrahydrofuran, dioxane, dimethoxyethane or diethyl ether, an alcohol such as methanol or ethanol, dimethyl sulfoxide, dimethylformamide, hexamethylphosphoric triamide, acetonitrile, and water, either singly or in combination with each other.

The reaction temperature is preferably 0 to 80°C, especially preferably 10 to 50 °C.

The reaction time varies depending upon the starting compound, the dehydrating agent and the reaction solvent used in the reaction. Usually, it is 10 minutes to 10 days, preferably 20 minutes to 5 days.

The 5-alkylidene-3-hydroxy-4-substituted cyclopentenone of formula (III) [including formula (III)-1] can be produced, for example, in accordance with the following reaction formula.

- 10 -



Specifically, it can be produced by reacting the corresponding 3-hydroxy-5-(1-hydroxy-aliphatic hydrocarbon)-4-substituted cyclopentanone derivative of formula (IV)' with methanesulfonyl chloride in the presence of a basic compound and then eliminating the protective group.

As can be understood from the above reaction formula, it is possible to subject the 5-alkylidene-3-hydroxy-4-substituted cyclopentanone of formula (III)' in which the 3-position hydroxyl group is protected by a t-butyldimethylsilyl group to a de-protection reaction using a deprotecting aid such as acetic acid thereby removing the protective group at the 3-position and forming the 5-alkylidene-3-hydroxy-4-substituted cyclopentanone in situ, and subsequently to subject this reaction mixture to the dehydration reaction in accordance with this invention.

According to the dehydration reaction of this invention, a corresponding compound (final desired compound) having a double bond formed between the 2- and 3-positions in formula (II) as a result of removal of the 3-position hydroxyl group is formed. When the resulting final compound has in the group W' and/or Y' a group removable by hydrolysis or deprotection or a group capable of forming a salt by salt-forming reaction and it is desired to obtain a final product by hydrolysis, deprotection or salt-forming reaction, the above dehydration reaction in the

- 11 -

process of this invention can be followed by such a reaction.

Groups capable of being removed by hydrolysis are, for example, carboacyl groups or ester groups.

- 5 The carboacyl groups can be hydrolyzed, for example, in an aqueous solution of sodium hydroxide, potassium hydroxide or calcium hydroxide, a water-alcohol mixture, a methanol or ethanol solution containing sodium methoxide, potassium methoxide or sodium
10 ethoxide. The ester groups can be hydrolyzed, for example, in water or a solvent containing water at a temperature of -10°C to $+60^{\circ}\text{C}$ for a period of about 10 minutes to about 24 hours using an enzyme such as lipase.

- 15 Groups capable of being removed by deprotection are, for example, groups forming an acetal linkage with the oxygen atom of the hydroxyl group, or tri(C_{1-7} hydrocarbon)silyl groups. The removal of the protective group can be performed suitably,
20 for example by using acetic acid, a pyridinium salt of p-toluenesulfonic acid, a cation exchange resin, etc. as a catalyst and water, tetrahydrofuran, diethyl ether, dioxane, acetone, acetonitrile, etc. as a reaction solvent when the protective group is a
25 group forming an acetal linkage together with the oxygen atom of the hydroxyl group. The reaction is carried out usually at a temperature of -78°C to $+30^{\circ}\text{C}$ for about 10 minutes to about 3 days. When the protective group is a tri(C_{1-7} hydrocarbon)-
30 silyl group, the deprotecting reaction may be carried out in the presence of acetic acid, tetrabutyl ammonium fluoride, cesium fluoride, etc. in the same reaction solvent as cited above at the same temperature and for the same period of time as mentioned
35 above.

When the final compound has a carboxyl group

- 12 -

in the molecule, it can then optionally be subjected to a salt-forming reaction to obtain the final compound as a carboxylate salt. The salt-forming reaction is known per se, and is carried out by neutralizing the carboxylic acid with a nearly equivalent of a basic compound such as sodium hydroxide, potassium hydroxide or sodium carbonate, or ammonia, trimethylamine, monoethanolamine or morpholine in a customary manner. The final desired compound can be isolated and purified, for example, by silica gel column chromatography, silica gel thin-layer chromatography, high-performance liquid chromatography, Florisil column chromatography, etc.


The 5-alkylidene-4-substituted-2-cyclopentenones of formula (I) in accordance with this invention can also be produced by treating the 5-(1-hydroxy-aliphatic hydrocarbon)-4-substituted-2-cyclopentenones of formula (II) with methane-sulfonyl chloride in the presence of basic compounds. The basic compounds may be amines such as dimethylaminopyridine, triethylamine, isopropylcyclohexylamine, isopropyldimethylamine and diisopropylamine. A reaction solvent may be used, and examples include halogenated hydrocarbons such as dichloromethane, chloroform, carbon tetrachloride, ethers such as diethyl ether and tetrahydrofuran, and aromatic hydrocarbons such as benzene and toluene. Methane-sulfonyl chloride is used generally in an amount of 1 to 10 moles per mole of the 5-(1-hydroxy-aliphatic hydrocarbon)-4-substituted-2-cyclopentenone of formula (II), and the reaction temperature may be from 0 to 50 °C, preferably 15 to 25 °C. The method disclosed in European Laid-Open Patent Publication No. 0079733 may be referred to in performing the above process.

In the second process of this invention, Wⁿ in

- 13 -

formula (IV) defining the starting material is an aliphatic hydrocarbon group having 1 to 12, and Y" is likewise an aliphatic hydrocarbon group having 1 to 12 carbon atoms. These aliphatic hydrocarbon groups may be substituted. Specific examples of W" and Y" may be the same as those given hereinabove for W' and Y'.


According to this invention, the second process is preferably carried out by producing a 4-substituted-5-(1-hydroxy-aliphatic hydrocarbon)-2-cyclopentenone represented by the following formula (II)-1



... (II)-1

wherein R¹, R², R³ and the symbol \equiv are the same as defined for formula (III)-1 above,

from a 5-(1-hydroxy-aliphatic hydrocarbon)-3-hydroxy-4-substituted cyclopentanone represented by the following formula (IV)-1



... (IV)-1

wherein R¹, R², R³, and the symbol \equiv are the same for formula (II)-1.

The dehydration reaction of the compound of formula (IV) or (IV)-1 can be carried out under quite

- 14 -

the same reaction conditions as in the aforesaid first process. In this reaction, the reaction temperature is preferably 0 to 130 °C, especially preferably 30 to 110 °C.

- 5 Deprotection, hydrolysis and salt-forming reaction can all be carried out in quite the same ways as described in regard to the first process.

Thus, according to this invention, there are provided 5-alkylidene-4-substituted-2-cyclopentenones
10 represented by the following formula (I)




wherein W and Y are as defined above,
and 5-(1-hydroxy-aliphatic hydrocarbon)-4-substituted-
2-cyclopentenones represented by the following
15 formula (II)



wherein W' and Y' are as defined above,
as novel 5-membered cyclic compounds.

Among the compounds of formula (I), those of
20 the following compounds (I)-1




wherein R¹ represents a hydrogen atom, or an

- 15 -

aliphatic hydrocarbon group having 1 to 10 carbon
atoms which may be substituted, R^2 represents a
hydrogen atom, or an aliphatic hydrocarbon group
having 1 to 9 carbon atoms which may be substituted,
5 R^3 represents a hydrogen atom, a hydroxyl group or
a protected hydroxyl group, and the symbol \equiv
represents a single, double or triple bond,
are preferred, and those of the following formula
(I)-2

10



wherein R^2 , R^3 , R^4 and the symbol \equiv
are as defined above, and the symbol \equiv
represents a single or double bond,
(A-type prostaglandins) are especially preferred.
15 R^1 in formulae (I)-1 and (II)-1 represents a
hydrogen atom or an aliphatic hydrocarbon group
having 1 to 10 carbon atoms. The aliphatic hydro-
carbon group may be substituted.

The aliphatic hydrocarbon group having 1 to 10
20 carbon atoms may be linear, branched or cyclic, and
may have a carbon-carbon double bond.

Examples of preferred aliphatic hydrocarbon
groups having 1 to 10 carbon atoms include linear or
branched alkyl or alkenyl groups and cycloalkyl
25 groups having 3 to 8 carbon atoms.

Specific examples of the alkyl groups having 1
to 10 carbon atoms include methyl, ethyl, n-propyl,
iso-propyl, n-butyl, sec-butyl, tert-butyl, n-pentyl,
n-hexyl, n-heptyl, n-octyl, n-nonyl and n-decyl.
30 Specific examples of the alkenyl groups having

- 16 -

1 to 10 carbon atoms include ethenyl, 1-propen-1-yl, 2-propen-1-yl, 3-buten-1-yl, 1-buten-1-yl, 4-penten-1-yl, 5-hexen-1-yl, 6-penten-1-yl, 7-octen-1-yl, 8-nonen-1-yl and 9-decen-1-yl. Of these, 1-propen-1-yl, 2-propen-1-yl, 3-buten-1-yl, 4-penten-1-yl and 5-hexen-1-yl are preferred.

Examples of the cycloalkyl groups having 3 to 8 carbon atoms are the same as those given for W' and Y' in formula (III).

10 R^2 in formulae (I)-1, (I)-2, (II)-1 and (II)-2 represents a hydrogen atom or an aliphatic hydrocarbon group having 1 to 9 carbon atoms. The aliphatic hydrocarbon group may be substituted.

Examples of the aliphatic hydrocarbon groups having 1 to 9 carbon atoms include linear or branched alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, t-butyl, n-pentyl, n-hexyl, 2-methyl-1-hexyl, 2-methyl-2-hexyl, n-heptyl and n-octyl, and the same C_{3-8} cycloalkyl groups as exemplified above.

20 R^3 in formulae (I)-1, (I)-2, (II)-1 and (II)-2 represents a hydrogen atom, a hydroxyl group or a protected hydroxyl group. Examples of the protective group for the hydroxyl group are tri(C_{1-7} hydrocarbon)silyl groups and groups forming an acetal linkage with the oxygen atom of the hydroxyl group.

Specific examples of preferred tri(C_{1-7} hydrocarbon)silyl groups include tri(C_{1-4} alkyl)silyl groups such as trimethylsilyl, triethylsilyl or t-butyldimethylsilyl, diphenyl(C_{1-4} alkyl)silyl groups such as t-butyldiphenylsilyl, and a tribenzylsilyl group.

Examples of the groups forming an acetal linkage together with the oxygen atom of the hydroxyl group include methoxymethyl, 1-ethoxyethyl, 2-methoxy-2-propyl, 2-ethoxy-2-propyl, (2-methoxyethoxy)methyl,

- 17 -

benzyloxymethyl, 2-tetrahydropyranyl, 2-tetrahydrofuranyl and 6,6-dimethyl-3-oxa-2-oxo-bicyclo[3.1.0]hex-4-yl groups. Of these, 2-tetrahydropyranyl, 2-tetrahydrofuranyl, 1-ethoxyethyl, 2-methoxy-2-propyl, (2-methoxyethoxy)methyl and 6,6-dimethyl-3-oxa-2-oxo-bicyclo[3.1.0]hex-4-yl groups are particularly preferred.

The following examples are given for the novel cyclic 5-membered compounds of the above formulae (I) [including formulae (I)-1 and (I)-2] and (II) [including formulae (II)-1 and (II)-2].

(i) Compounds of formula (I)

- (100) 4-Butyl-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (102) 4-octyl-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (104) 4-(1-octenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (106) 4-(3-hydroxy-1-octenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (108) 4-(3-hydroxy-3-cyclohexyl-1-propenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (110) 4-(3-hydroxy-3-cyclopentyl-1-propenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (112) 4-(2-propenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (114) 4-(3-hydroxy-5-methyl-nonenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (116) 4-(3-hydroxy-3-phenyl-1-propenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (118) 4-(1-hydroxy-3-phenyl-1-propenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (120) 4-(3-hydroxy-3-methyl-1-propenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (122) 4-(3-hydroxy-5,5-dimethyl-1-octenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,

- 18 -

- (124) 4-(3-hydroxy-4-phenoxy-1-butenyl)-5-(6-carboxyhexylidene)-2-cyclopentenone,
(126) 4-butyl-5-(6-carboxy-2-hexenylidene)-2-cyclopentenone,
5 (128) 4-butyl-5-(6-carboxy-2-hexynylidene)-2-cyclopentenone,
(130) 4-butyl-5-(6-carboxy-5-hexenylidene)-2-cyclopentenone,
(132) 4-octenyl-5-(6-carboxy-2-hexenylidene)-
10 2-cyclopentenone,
(134) 4-octenyl-5-(6-carboxy-2-hexenylidene)-2-cyclopentenone,
(136) 4-(3-hydroxy-1-octenyl)-5-(6-carboxy-2-hexynylidene)-2-cyclopentenone,
15 (138) 4-(3-hydroxy-3-cyclopentyl-1-propenyl)-5-(6-carboxy-2-hexenylidene)-2-cyclopentenone,
(140) 4-(3-hydroxy-3-cyclohexyl-1-propenyl)-5-(6-carboxy-2-hexynylidene)-2-cyclopentenone,
(142) 4-butyl-5-(2-methylpropylidene)-2-cyclopentenone,
20 (144) 4-butyl-5-(2,2-dimethylpropylidene)-2-cyclopentenone,
(146) 4-(3-hydroxy-1-octenyl)-5-butylidene-2-cyclopentenone,
25 (148) 4-butyl-5-(3-phenyl-2-propenylidene)-2-cyclopentenone,
(150) 4-octyl-5-(2-methylpropylidene)-2-cyclopentenone,
(152) 4-(1-octenyl)-5-(6-carboxy-2-hexenylidene)-2-cyclopentenone,
30 (154) 4-butyl-5-heptylidene-2-cyclopentenone,
(156) 4-octyl-5-heptylidene-2-cyclopentenone,
(158) 4-(3-hydroxy-1-octenyl)-5-heptylidene-2-cyclopentenone,
35 (160) 4-butyl-5-(7-hydroxyheptylidene)-2-cyclopentenone,

- 19 -

- (162) 4-octyl-5-(7-hydroxyheptylidene)-2-cyclopentenone,
- (164) 4-(3-hydroxy-1-octenyl)-5-(7-hydroxyheptylidene)-2-cyclopentenone,
- 5 (166) 4-(1-octenyl)-5-(7-hydroxyheptylidene)-2-cyclopentenone,
- (168) 4-(3-hydroxy-4-m-fluorophenoxy)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (170) 4-(3-hydroxy-4-m-trifluoromethylphenyl)-
- 10 5-(6-carboxyhexylidene)-2-cyclopentenone,
- (172) 4-(1-octyne)-5-(6-carboxyhexylidene)-2-cyclopentenone,
- (174) methyl ester of (100),
- (176) methyl ester of (102),
- 15 (178) methyl ester of (104),
- (180) methyl ester of (106),
- (182) ethyl ester of (108),
- (184) ethyl ester of (110),
- (186) ethyl ester of (112),
- 20 (188) propyl ester of (114).
- (190) propyl ester of (120),
- (192) sodium salt of (124),
- (194) sodium salt of (126),
- (196) sodium salt of (128),
- 25 (198) aluminum salt of (130),
- (200) aluminum salt of (136),
- (202) aluminum salt of (138),
- (204) aluminum salt of (140),
- (ii) Compound of formula II
- 30 (300) 4-butyl-5-(1-hydroxybutyl)-2-cyclopentenone,
- (302) 4-butyl-5-(1-hydroxy-1-methylpropyl)-2-cyclopentenone,
- (304) 4-butyl-5-(1-hydroxy-2,2-dimethylpropyl)-
- 35 2-cyclopentenone,
- (306) 4-butyl-5-(1-hydroxy-3-phenyl-2-propen-

- 20 -

- 1-yl)-2-cyclopentenone,
(308) 4-butyl-5-(1-hydroxybutyl)-2-cyclopentenone,
(310) 4-octyl-5-(1-hydroxybutyl)-2-cyclopentenone,
5 none
(312) 4-(2-propenyl)-5-(1-hydroxybutyl)-2-cyclopentenone,
(314) 4-(1-octenyl)-5-(1-hydroxybutyl)-2-cyclopentenone,
10 (316) 4-(3-hydroxy-1-octenyl)-5-(1-hydroxybutyl)-2-cyclopentenone,
(318) 4-butyl-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
(320) 4-octyl-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
15 (322) 4-(1-octenyl)-5-(1-hydroxy-6-carboxyhexyl)-1-cyclopentenone,
(324) 4-decyl-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
20 (326) 4-(5-methyl-1-nonenyl)-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
(328) 4-(3-hydroxy-5-methyl-1-nonenyl)-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
(330) 4-(3-cyclohexyl-1-propenyl)-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
25 (332) 4-(1-octenyl)-5-(1-hydroxy-8-carboxyoctyl)-2-cyclopentenone,
(334) 4-(1-octenyl)-5-(1-hydroxy-5-carboxypentyl)-2-cyclopentenone,
30 (336) 4-(3-hydroxy-1-octenyl)-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
(338) 4-(3-hydroxy-1-octenyl)-5-(1-hydroxy-6-carboxy-5-hexene-1-yl)-2-cyclopentenone,
(340) 4-(3-hydroxy-1-octenyl)-5-(1-hydroxy-6-carboxy-3-hexene-1-yl)-2-cyclopentenone,
35 (342) 4-(3-hydroxy-1-octenyl)-5-(1-hydroxy-6-

- 21 -

- carboxy-2-hexene-1-yl)-2-cyclopentenone,
 (344) 4-(3-hydroxy-5-methyl-1-nonenyl)-5-(1-hydroxy-6-carboxy-5-hexene-1-yl)-2-cyclopentenone,
 (346) 4-(3-hydroxy-4-methyl-1-nonenyl)-5-(1-hydroxy-6-carboxy-2-hexyne-1-yl)-2-cyclopentenone,
5 (348) 4-(3-hydroxy-3-cyclohexyl-1-propenyl)-5-(1-hydroxy-6-carboxy-2-hexyne-1-yl)-2-cyclopentenone,
 (350) 4-(3-hydroxy-3-cyclohexyl-1-propenyl)-5-(1-hydroxy-8-carboxy-2-octyne-1-yl)-2-cyclopentenone,
10 (352) 4-(3-hydroxy-4-cyclohexyl-1-butenyl)-5-(1-hydroxy-6-carboxy-4-hexene-1-yl)-2-cyclopentenone,
 (354) 4-(3-hydroxy-3-cyclopentyl-1-propenyl)-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
 (356) 4-(3-hydroxy-3-phenoxy-1-propenyl)-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
15 (358) 4-(3-hydroxy-4-phenoxy-1-butene)-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
 (360) 4-(3-hydroxy-4,4-dimethyl-1-octenyl)-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
20 (362) 4-(3-hydroxy-5,5-dimethyl-1-octenyl)-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone,
 (364) methyl ester of (300),
 (366) methyl ester of (302),
 (368) methyl ester of (304),
25 (370) methyl ester of (306),
 (372) ethyl ester of (308),
 (374) ethyl ester of (310),
 (376) ethyl ester of (314).
 (378) propyl ester of (320),
30 (380) propyl ester of (324),
 (382) sodium salt of (330),
 (384) sodium salt of (334),
 (386) sodium salt of (350),
 (388) aluminum salt of (352).
35 (390) aluminum salt of (356),
 (392) aluminum salt of (358).

- 22 -

Investigations of the present inventors have shown that the novel cyclic 5-membered compounds of this invention, i.e. the 5-alkylidene-4-substituted 2-cyclopentenones of formula (I) and the 5-(alpha-
5 hydroxy-aliphatic hydrocarbon)-4-substituted-2-cyclopentenones of formula (II), and the 5-alkylidene-3-hydroxy-4-substituted cyclopentanones of formula (III) used as starting materials for the production of the compounds of formula (I) in the first process
10 of this invention have excellent pharmacological activities.

Accordingly, the present invention also provides a pharmaceutical composition comprising as an active ingredient a 5-membered cyclic compound selected from the group consisting of the 5-alkylidene-
15 4-substituted-2-cyclopentenones of formula (I), the 5-(1-hydroxy-hydrocarbon)-4-substituted-2-cyclopentenones of formula (II), and the 5-alkylidene-3-hydroxy-4-substituted cyclopentanones of formula (III), and a
20 pharmaceutically acceptable carrier.

The 5-membered cyclic compounds of formulae (I), (II) and (III) in accordance with this invention exhibit strong anticancer activity in low concentrations against L1210 leukemia cells and are very
25 useful as antitumor agents. They are also useful as compounds having antiviral activity.

According to this invention, the 5-membered cyclic compounds can be administered orally, or parenterally through intrarectal, subcutaneous,
30 intramuscular and intravenous routes, for example. For oral administration, the compounds of this invention may be formulated into solid or liquid preparations. Examples of the solid preparations are tablets, pills, powders and granules. In these solid
35 preparations, at least one of the 5-membered cyclic compounds is mixed with sodium bicarbonate, calcium

- 23 -

carbonate, potato starch, sucrose, mannitol, carboxy-methyl cellulose, etc. These preparations can be formed in accordance with customary operations. The solid preparations may also include a lubricant, a
5 sweetener, a stabilizer, an antiseptic, etc. such as calcium stearate, magnesium stearate or glycerol.

Examples of the liquid preparations for oral administration are emulsions, solutions, suspensions, syrups, and elixirs. The liquid preparations may
10 further include a wetting agent, a suspending aid, a sweetener, a flavor, an aroma, a stabilizer, etc. The liquid preparations may be filled in capsules made of an absorbable material such as gelatin.

For intrarectal administration, ordinary
15 suppositories such as soft gelatin capsules are used.

Examples of preparations for parenteral administration through other routes are preparations for subcutaneous, intramuscular or intravenous injection in the form of aseptic aqueous or non-aqueous
20 solutions, suspensions and emulsions. The non-aqueous solutions and suspensions may include propylene glycol, polyethylene glycol, olive oil or injectable organic esters such as ethyl oleate. Such preparations may also contain an antiseptic, an
25 emulsifier, a dispersant, a stabilizer, etc. These injectable preparations can be made aseptic by filtration through a bacteria-holding filter, blending of a germicide, or irradiation.

The dose of the 5-membered cyclic compound of
30 this invention differs depending upon the condition, age, sex and body weight of a subject to which it is to be administered, the route of administration, etc. Usually, it can be administered in a dose of about $1\mu\text{g}$ to 100 mg/kg-body weight/day. The dose may be a
35 single dose, or may be divided into several portions, for example 2 to 6 portions.

- 24 -

The pharmaceutical composition of this invention is preferably used as a medicament in unit dosage form.

The following Examples illustrate the present invention in greater detail.

5 Example 1

Synthesis of (7E)-7,8-dehydro PGE₁:

(1) 450 mg (0.75 mmole) of 7-hydroxy PGE₁ 11,15-bis(t-butyldimethylsilyl)ether was dissolved in 7 ml of anhydrous dichloromethane, and 367 mg (30 mmoles) of dimethylaminopyridine was added. Methanesulfonyl chloride (116 microliters; 1.5 mmoles) was added, and the mixture was stirred at room temperature for 16 hours. A saturated aqueous solution of sodium chloride was added, and the mixture was adjusted to pH 2 with oxalic acid and extracted with dichloromethane. The extract was dried over anhydrous magnesium sulfate, filtered, concentrated and subjected to silica gel column chromatography (silica gel 20 g; eluent, hexane:acetone=20:1 ~ 5:1) to give 72 mg (yield 16%) of (7E)-7,8-dehydro PGE₁ 11,15-bis(t-butylmethylsilyl)ether and 17 mg (yield 4%) of (7Z)-7,8-dehydro PGE₁ 11,15-bis(t-butyldimethylsilyl)ether.

25 Spectral data of (7E)-7,8-dehydro PGE₁ 11,15-bis(t-butyldimethylsilyl)ether:-

TLC: R_f=0.43 (hexane:acetone=2:1)

IR (liquid film):

3600-2400, 1713, 1650, 1461, 1255, 1073, 833
772, 929 cm⁻¹.

30 NMR (CDCl₃) δ:

0-0.2 (m, 12H), 0.84 (s, 9H), 0.87 (s, 9H),
0.7-1.1 (m, 3H), 1.0-3.0 (m, 20H), 3.2-3.7
(m, 1H), 3.8-4.3 (m, 2H), 5.3-5.7 (m, 2H),
6.66 (dt, 1H, J=7.5, 2.0Hz), 9.0-9.8 (m, 1H).

35 Spectral data of (7E)-7,8-dehydro PGE₁ 11,15-bis(t-butyldimethylsilyl)ether:-

- 25 -

TLC: Rf=0.52 (hexane:acetone=2:1)

IR (liquid film):

3600-2400, 1740, 1648, 1460, 1254, 1077,
836, 773, 757 cm^{-1} .5 NMR (CDCl_3) δ :0-0.2 (m, 12H), 0.88 (s, 9H), 0.90 (s, (H),
0.7-1.1 (m, 3H), 1.0-3.0 (m, 20H), 3.3-3.5
(m, 1H), 3.8-4.3 (m, 2H), 4.8-6.0 (m, 1H),
5.4-5.8 (m, 2H)

10 (2) 22 mg (38 micromoles) of (7E)-7,8-dehydro
PGE₁ 11,15-bis(t-butyldimethylsilyl)ether was added
to 1.0 ml of a hydrogen fluoride-acetonitrile so-
lution (prepared by adding 0.5 ml of 47% hydrofluoric
acid to 10 ml of acetonitrile), and the mixture was
15 stirred at room temperature for 20 minutes. A satu-
rated aqueous solution of sodium bicarbonate was
added, and the mixture was acidified with oxalic acid
and extracted with ethyl acetate. The organic layer
was washed with a saturated aqueous solution of
20 sodium chloride, and dried over anhydrous magnesium
sulfate. The dried product was filtered, concen-
trated and subjected to silica gel thin-layer chromato-
graphy (developing solvent, hexane:acetone:acetic
acid=1:1:0.01) to give 3.7 mg (yield 28%) of (7E)-7,8-
25 dehydro PGE₁.

Spectral data of (7E)-7,8-dehydro PGE₁:-

TLC: Rf=0.47 (hexane:acetone=1:2)

IR (CHCl_3 solution):3400, 1715, 1642, 973 cm^{-1} .30 NMR (CDCl_3) δ :0.88 (brt, 3H), 1.0-1.9 (m, 14H),
1.9-2.9 (m, 5H), 2.9-3.9 (m, 5H),
3.9-4.5 (m, 2H), 5.3-5.9 (m, 2H),
6.7-7.0 (m, 1H).35 Example 2Synthesis of (7E)-7,8-dehydro PGA₁:

- 26 -

160 mg of (7E)-7,8-dehydroprostaglandin E_1 was dissolved in 3 ml of tetrahydrofuran, and 2 ml of 0.5N hydrochloric acid was added. The mixture was stirred at room temperature for 4 days. A saturated aqueous solution of sodium chloride was added, and the mixture was extracted with ethyl acetate. The organic layers were combined, washed with a saturated aqueous solution of sodium chloride and dried over anhydrous magnesium sulfate. The dried product was filtered, concentrated and purified by silica gel thin-layer chromatography to give 95 mg (yield 63%) of (7E)-7,8-dehydroprostaglandin A_1 .

TLC: R_f =0.58 (developing solvent,
hexane:acetone=1:2)

IR ($CHCl_3$ solution):
3650-2400, 1698, 1646. 967 cm^{-1} .

NMR ($CDCl_3$) δ :
0.88 (t, 3H, J =6.0Hz), 1.0-2.0 (m, 14H),
2.0-2.8 (m, 4H), 3.4-4.6 (m, 4H), 5.1-5.9
(m, 2H), 6.34 (dd, 1H, J =6.0, 2.4Hz),
6.61 (dd, 1H, J =7.8, 1.5Hz), 7.36 (dd, 1H,
 J =6.0, 2.7Hz).

Example 3

Synthesis of (7E)-7,8-dehydro PGA_1 :

22 mg of (7E)-7,8-dehydroprostaglandin E_1 , 11,15-bis(*t*-butylmethylsilyl)ether was dissolved in a solution containing 1 ml of acetonitrile and 50 microliters of 47% hydrofluoric acid, and the mixture was stirred at room temperature for 20 minutes. A saturated aqueous solution of sodium bicarbonate was added, and the mixture was adjusted to pH 1 with oxalic acid, and extracted with ethyl acetate. The organic layers were combined, washed with a saturated aqueous solution of sodium chloride, and dried over anhydrous magnesium sulfate. The dried product was

- 27 -

filtered, concentrated and purified by silica gel thin-layer chromatography to give 4.7 mg (yield 37%) of (7E)-7,8-dehydroprostaglandin A₁.

Example 4

- 5 Synthesis of (7E)-7,8-dehydro PGA₁ methyl ester and 12-epi-(7E)-7,8-dehydro PGA₁ methyl ester:-

- 1.0 mg of a mixture of (7E)-7,8-dehydro PGE₁ methyl ester 11,15-bis(t-butyldimethylsilyl)ether and
10 15-epi-ent-(7E)-7,8-dehydro PGE₁ methyl 11,15-bis-(t-butyldimethylsilyl)ether was suspended in a mixed solvent consisting of acetic acid, tetrahydrofuran and water in a ratio of 2:1:1, and the suspension was stirred at 50° for 14 hours and then at 60°C for 3
15 hours. After concentration, water and sodium carbonate were added to the residue to perform neutralization. The mixture was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous
20 magnesium sulfate, filtered, concentrated, and subjected to silica gel column chromatography (silica gel 200 g; hexane:ethyl acetate=1:1) to give 184 mg (yield 1%) of 12-epi-(7E)-7,8-dehydro PGA₁ methyl ester and 184 mg (yield 31%) of (7E)-7,8-dehydro
25 PGA₁ methyl ester.

Spectral data of 12-epi-(7E)-7,8-dehydro PGA₁ methyl ester:-

TLC: R_f=0.57 (hexane:ethyl acetate=1:3)

IR (liquid film):

- 30 3560, 1738, 1700, 1648, 1577 cm⁻¹.

NMR (CDCl₃)δ:

- 0.90 (brt, 3H), 1.0-2.0 (m, 14H), 2.0-2.7
(m, 5H), 3.66 (s, 3H), 3.75-4.45 (m, 2H),
5.45 (dd, 1H, J=15.5, 6.5Hz), 5.71 (dd, 1H,
35 J=15.5, 6.0Hz)m, 6.30 (dd, 1H, J=6.0, 2.0Hz),
6.63 (brt, 1H, J=8.0Hz), 7.41 (dd, 1H,

- 28 -

$J=6.0, 3.0\text{Hz}$).

Spectral data of (7E)-7,8-dehydro PGA₁ methyl ester:-

TLC: $R_f=0.51$ (hexane:ethyl acetate=1:3)

5 IR (liquid film):

3450, 1739, 1701, 1648, 1578 cm^{-1} .

NMR (CDCl_3) δ :

0.90 (brt, 3H), 1.0-2.0 (m, 14H), 2.0-2.8 (m, 5H), 3.66 (s, 3H), 3.7-4.5 (m, 2H),
10 5.46 (dd, 1H, $J=15.5, 6.5\text{Hz}$), 5.71 (dd, 1H, $J=15.5, 6.0\text{Hz}$), 6.34 (dd, 1H, $J=6.0, 2.0\text{Hz}$), 6.63 (brt, 1H, $J=8.0\text{Hz}$), 7.41 (dd, 1H, $J=6.0, 3.0\text{Hz}$).

Example 5

15 Synthesis of 4-(1-octenyl)-5-(6-methoxycarbonylhexylidene)-2-cyclopentenone:-

(1) 3.1 g (6.4 millimoles) of 4-(1-octenyl)-5-(1-hydroxy-6-methoxycarbonylhexyl)-3-(t-butyl-dimethylsilyloxy)cyclopentanone was dissolved in
20 40 ml of dichloromethane, and 3.92 g (32.1 mmols) of dimethylaminopyridine was added. With ice cooling and stirring, 1.0 ml (12.9 mmols) of methanesulfonyl chloride was added. The mixture was stirred at 0°C for 5 minutes, and then at room temperature for
25 12 hours. Furthermore, 0.78 g (6.4 millimoles) of dimethylaminopyridine was added, and the mixture was stirred for 100 minutes. The mixture was poured into 20 ml of 0.5N hydrochloric acid and washed. The aqueous layer was extracted with dichloromethane.
30 The organic layers were combined, washed first with a saturated aqueous solution of sodium bicarbonate and then with a saturated aqueous solution of sodium chloride, and dried over anhydrous sodium sulfate. The dried product was filtered, concentrated and
35 subjected to silica gel column chromatography (silica gel 150 g; eluent, hexane:ethyl acetate=20:1 - 7:1)

- 29 -

to give 1.83 g (yield 61%) of 4-(1-octenyl)-5-(6-methoxycarbonylhexylidene)-3-(t-butyldimethylsilyloxy)-cyclopentanone. The spectral data of this compound were as follows.-

5 TLC: Rf=0.45 (hexane:ethyl acetate=5:1)

NMR (CDCl₃) δ :

0-0.2 (m, 6H), 0.83 (s, 9H), 0.7-1.1

(m, 3H), 1.0-2.8 (m, 22H), 3.1-3.4 (m, 1H),

3.59 (s, 3H), 3.9-4.3 (m, 1H), 5.1-5.5

10 (m, 2H), 6.61 (td, 1H, J=7.5, 2.0Hz).

(2) 1.3 g (2.8 mmoles) of 4-(1-octenyl)-5-(6-methoxycarbonylhexylidene)-3-(t-butyldimethylsilyloxy)-cyclopentanone was dissolved in 40 ml of a solvent consisting of acetic acid, tetrahydrofuran and water in a ratio of 2:1:1, and the solution was stirred at 15 60°C for 15 hours. Toluene was added, and the mixture was concentrated under reduced pressure. A saturated aqueous solution of sodium bicarbonate was add. The mixture was extracted with ethyl acetate 20 thre times. The organic layers were combined, washed with a saturated aqueous solution of sodium chloride and dried over anhydrous magnesium sulfate. The dried product was filtered, concentrated and subjected to silica gel column chromatography (silica 25 gel 60 g; eluent, hexane:ethyl acetate=7:1 - 1:1) to give 32 mg (yield 3%) of 4-(1-octenyl)-5-(6-methoxycarbonylhexylidene)-2-cyclopentanone, a less polar isomer, and 635 mg (yield 70%) of 4-(1-octenyl)-5-(6-methoxycarbonylhexylidene)-2-cyclopentenone, a more 30 polar isomer.

Spectral data of the less polar isomer:-

TLC: Rf=0.33 (hexane:ethyl acetate=5:1)

NMR (CDCl₃) δ :

0.85 (brt, 3H, J=4.7Hz), 1.0-2.5 (m, 18H),

35 2.5-3.0 (m, 2H), 3.58 (s, 3H), 3.5-4.0

(m, 1H), 5.10 (dd, 1H, J=15.6, 8.0Hz),

- 30 -

5.48 (dd, 1H, J=15.6, 6.0Hz), 5.88 (brt, 1H, J=7.2Hz), 6.16 (dd, 1H, J=6.0, 2.2Hz), 7.18 (dd, 1H, J=6.0, 2.4Hz).

Spectral data of the more polar isomer:-

5 TLC: Rf=0.25 (hexane:ethyl acetate=5:1)

NMR (CDCl₃) δ :

0.85 (brt, 3H, J=4.2Hz), 1.0-2.5 (m, 20H),
3.58 (s, 3H), 3.7-4.1 (m, 1H), 5.12 (dd, 1H, J=15.0, 7.7Hz)m, 5.52 (dt, 1H, J=15.0, 6.2Hz), 6.19 (dd, 1H, J=5.8, 1.0Hz),
10 6.49 (brt, 1H, J=7.8Hz), 7.24 (dd, 1H, J=6.2, 2.2Hz).

Example 6

15 Synthesis of 4-butyl-5-(6-methoxycarbonyl-hexylidene)-3-hydroxycyclopentanone:-

By using 4-butyl-5-(6-methoxycarbonyl-1-hydroxy-hexyl)-3-t-butyldimethylsilyloxycyclopentanone, 4-butyl-5-(6-methoxycarbonylhexylidene)-3-hydroxycyclopentanone was prepared in the same way as in
20 Example 1. The resulting product had the following spectral data.

IR (liquid film):

3450, 1735, 1718, 1641 cm⁻¹.

NMR (CDCl₃) δ :

25 0.89 (brt, 3H, J=5.0Hz), 1.0-2.7 (m, 18H), 2.7-3.3 (m, 2H), 3.64 (s, 3H), 4.0-4.3 (m, 1H), 6.68 (dt, 1H, J=7.5, 2.0Hz).

Example 7

30 Synthesis of 4-butyl-5-(6-methoxycarbonyl-hexylidene)-2-cyclopentenone:-

81 mg (0.28 mmole) of 4-butyl-5-(6-methoxycarbonylhexylidene)-3-hydroxycyclopentanone obtained in Example 6 was dissolved in a mixture consisting of 2 ml of acetic acid, 1 ml of tetrahydrofuran and 1 ml of water, and the solution was stirred at 60°C for
35 7 hours and then at 90°C for 50 hours. After the

- 31 -

reaction, the reaction mixture was concentrated under reduced pressure and neutralized with an aqueous solution of sodium bicarbonate. The mixture was extracted with ethyl acetate. The organic layer
5 separated was washed with a saturated aqueous solution of sodium chloride, and dried over anhydrous magnesium sulfate. The dried product was filtered, concentrated and purified by silica gel column chromatography (silica gel 10 g; hexane:ethyl acetate=6:1) to
10 give 61 mg (0.22 mmole; 81%) of 4-butyl-5-(6-methoxycarbonylhexylidene)-2-cyclopentenone. The resulting compound had the following spectral data.

IR (liquid film):

1739, 1703, 1656, 1580 cm^{-1} .

15 NMR (CDCl_3 : δ (ppm)):

0.89 (3H, t), 1.0-2.0 (12H, m), 2.0-2.6

(4H, m), 3.3-3.8 (1H, m), 3.67 (3H, s)

6.35 (1H, dd, $J=6.0, 2.0\text{Hz}$), 6.56 (1H, t),

7.59 (1H, dd, $J=6.0, 3.0\text{Hz}$).

20 Example 8

Synthesis of 4-octyl-5-(6-carboxyhexylidene)-3-hydroxycyclopentanone:-

By using 4-octyl-5-(6-carboxyl-1-hydroxyhexyl)-3-t-butyltrimethylsilyloxycyclopentanone, 4-octyl-5-
25 (6-carboxyhexylidene)-3-hydroxycyclopentanone was prepared in the same way as in Example 1. The resulting compound had the following spectral data.

IR (liquid film):

3430, 1738, 1720, 1638 cm^{-1} .

30 NMR (CDCl_3) δ :

0.84 (brt, 3H, $J=4.7\text{Hz}$), 1.0-2.6 (m, 22H),

3.2-3.7 (m, 2H), 3.57 (s, 3H), 4.0-4.3

(m, 1H), 5.2-5.5 (m, 2H), 6.62 (dd, 1H,

$J=7.4, 2.0\text{Hz}$)

35

- 32 -

Example 9

Synthesis of 4-octyl-5-(6-carboxyhexylidene)-2-cyclopentenone:-

169 mg (0.5 mmole) of 4-octyl-5-(6-carboxyhexyl-
5 idene)-3-hydroxycyclopentanone was dissolved in 3 ml
of tetrahydrofuran, and 1 ml of 0.5N hydrochloric acid
was added. The mixture was stirred at room tempera-
ture for 4 days. A saturated aqueous solution of
sodium chloride was added, and the mixture was ex-
10 tracted with ethyl acetate. The organic layer was
washed with a saturated aqueous solution of sodium
chloride, and dried over anhydrous magnesium sulfate.
The dried product was filtered, concentrated, and
purified by preparative silica gel-thin-layer chromato-
15 graphy to give 12 mg (0.35 mmole; yield 70%) of
4-octyl-5-(6-carboxyhexylidene)-2-cyclopentanone.
The resulting compound had the following spectral
data.

IR (CHCl_3 solution):

20 3650-2400, 1700, 1646, 1580 cm^{-1} .

NMR (CDCl_3 : δ (ppm)):

0.89 (t, 3H), 1.0-2.0 (m, 20H), 2.0-2.6
(m, 4H), 3.3-3.8 (m, 1H), 6.33 (dd, 1H,
J=6.2Hz), 6.60 (t, 1H), 7.53 (dd, 1H,
25 J=6.3Hz), 11.5 (1H).

Example 10

Synthesis of 4-butyl-5-(6-methoxycarbonyl)-2-hexynylidene)-3-hydroxycyclopentanone:-

By using 4-butyl-5-(6-methoxycarbonyl-1-hydroxy-
30 2-hexynyl)-3-t-butyldimethylsilyloxycyclopentanone,
4-butyl-5-(6-methoxycarbonyl-2-hexynylidene)-3-hydroxy-
cyclopentanone was prepared in the same way as in
Example 1. The resulting compound had the following
spectral data.

35 IR (liquid film):

3440, 2200, 1735, 1715, 1608 cm^{-1} .

- 33 -

NMR (CDCl_3) δ :

0.85 (brt, 3H, $J=4.5\text{Hz}$), 1.0-2.7 (m, 14H),
2.7-3.3 (m, 2H), 3.64 (s, 3H), 4.0-4.3
(m, 1H), 6.3-6.5 (m, 1H)

5 Example 11

Synthesis of 4-butyl-5-(6-methoxycarbonyl-
2-hexynylidene)-2-cyclopentenone:-

29 mg (0.1 mmole) of 4-butyl-5-(6-methoxy-
carbonyl-2-hexynylidene)-3-hydroxycyclopentanone was
10 dissolved in a mixture of 1 ml of acetic acid, 0.5 ml
of tetrahydrofuran and 0.5 ml of water and the solu-
tion was stirred at 50°C for 18 hours in the same
way as in Example 1. The reaction mixture was worked
up and separated in the same wy as in Example 7 to
15 give 13 mg (0.047 mmole; 47% yield) of 4-butyl-5-(6-
methoxycarbonyl-2-hexynylidene)-2-cyclopentenone.
The resulting compound had the following spectral
data.

IR (liquid film):

20 2190, 1730, 1680, 1620, 1565 cm^{-1} .NMR (CDCl_3 : δ (ppm)):

0.88 (3H, t), 1.0-2.2 (8H, m), 2.2-2.8
(4H, m), 3.3-3.8 (1H, m), 3.67 (3H, s),
6.40 (1H, dd, $J=6.0, 2.0\text{Hz}$), 6.55 (1H),
25 7.53 (1H, dd, $J=6.0, 3.0\text{Hz}$).

Example 12

Synthesis of 4-(3-t-butyl dimethylsilyloxy-
3-cyclopentyl-1-propenyl)-5-(6-methoxy-
carbonyl-2-hexynylidene)-3-t-butyl dimethyl-
30 silyloxycyclopentanone:-

By using 4-(3-t-butyl dimethylsilyloxy)-3-
cyclopentyl-1-propenyl-5-(6-methoxycarbonyl-1-
hydroxy-2-hexynylidene)-3-t-butyl dimethylsilyloxy-
cyclopentanone, 4-(3-t-butyl dimethylsilyloxy-3-
35 cyclopentyl-1-propenyl)-5-(6-methoxycarbonyl-2-
hexylidene)-3-t-butyl dimethylsilyloxycyclopentanone

- 34 -

was prepared in the same way as in Example 1. The spectral data of the resulting compound were as follows:-

IR:

5 2218, 1749, 1730, 1620, 1254, 833, 771 cm^{-1} .

NMR (CDCl_3 , δ (ppm)):

0-0.2 (m, 12H), 0.89 (s, 18H), 1.1-2.5
(m, 17H), 3.3-3.65 (m, 1H), 3.67 (s, 3H),
3.75-4.5 (m, 2H), 5.4-5.9 (m, 2H), 6.4-6.7
10 (m, 1H).

Example 13

Synthesis of 4-(3-hydroxy-3-cyclopentyl-
1-propenyl)-5-(6-methoxycarbonyl-2-hexynyl-
iodene)-2-cyclopentenone:-

15 48 mg (81 micromoles) of 3-t-butyldimethyl-
silyloxy-4-(3-t-butyldimethylsilyloxy-3-cyclopentyl-
1-propenyl)-5-(6-methoxycarbonyl-2-hexylidene)cyclo-
pentanone was dissolved in a mixture of 1.5 ml of
acetic acid, 1 ml of tetrahydrofuran and 1 ml of
20 water, and the solution was stirred at 40°C for
35 hours. [In the early stage of the reaction,
3-hydroxy-4-(3-hydroxy-3-cyclopentyl-1-propenyl)-
5-(6-methoxycarbonyl-2-hexynylidene)cyclopentanone
formed. This was confirmed from the fact that it
25 agreed in thin-layer chromatography with a separately
prepared sample.] After the reaction, the reaction
mixture was concentrated under reduced pressure and
neutralized with sodium bicarbonate. The mixture was
extracted with ethyl acetate. The organic layer
30 separated with a saturated aqueous solution of sodium
chloride, and dried over anhydrous magnesium sulfate.
The dried product was filtered, concentrated, and
purified by silica gel column chromatography (silica
gel 3 g; hexane:ethyl acetate=3:1) to give 19 mg (55
35 micromoles; yield 69%) of 4-(3-hydroxy-3-cyclopentyl-
1-propenyl)-5-(6-methoxycarbonyl-2-hexynylidene-2-

- 35 -

cyclopentanone. The resulting compound had the following spectral data.

IR (liquid film):

3450, 2190, 1728, 1683, 1620, 1565,
743 cm^{-1} .

NMR (CDCl_3 : δ (ppm)):

0.9-2.3 (m, 11H), 2.3-2.8 (m, 5H), 3.70
(s, 3H), 3.45-4.35 (m, 2H), 5.5-5.9 (m, 2H),
6.3-6.7 (m, 2H), 7.4-7.7 (m, 1H).

10 Example 14

Synthesis of 7(E)-7,8-dehydro-17(S),20-dimethylprostaglandin A_1 methyl ester:-

In the same way as in Example 13, 7(E)-7,8-dehydro-17(S),20-dimethylprostaglandin A_1 methyl ester was obtained from 7(E)-7,8-dehydro-17(S),20-dimethylprostaglandin E_1 methyl ester-11,15-bis-t-butyldimethylsilyl ester.

NMR (CDCl_3 : δ (ppm)):

0.8-1.0 (m, 6H), 1.0-2.0 (m, 15H), 2.0-2.6
(m, 4H), 3.65 (s, 3H), 4.15 (m, 3H), 5.4
(dd, 1H, $J=15.6\text{Hz}$), 5.75 (dd, 1H, $J=15.6\text{Hz}$),
6.35 (dd, 1H, $J=6.2\text{Hz}$), 6.6 (t, 1H, $J=6\text{Hz}$),
7.4 (dd, 1H, $J=6.2\text{Hz}$).

Example 15

25 Synthesis of 3-n-butyl-4-t-butyldimethylsilyloxy-2-(1-hydroxybutyl)cyclopentanone and 4-n-butyl-5-(1-hydroxybutyl)-2-cyclopentenone:-

(1) 390 mg (2.05 mmoles) of cuprous iodide was
30 taken into reaction vessel. The reaction vessel was purged with argon, and 20 ml of dry ether and 1.02 ml (4.1 mmoles) of tributyl phosphine were added. The mixture was stirred for 10 minutes. The mixture was cooled to -78°C , and 1.25 ml (0.05 mmoles) of
35 n-butyllithium (as a 1.64M hexane solution) was added, and the mixture was stirred for 5 minutes.

- 36 -

The solution of 425 mg (2.0 mmoles) of 4-t-butyldi-methylsilyloxy-2-cyclopentenone in 5 ml of dry ether was added to the mixture, and the mixture was stirred at -78°C for 10 minutes and then at -40°C for 5 20 minutes. A solution of 148 mg (2.05 mmoles) of butanal in 5 ml of dry ether was added, and the mixture was stirred at -40°C for 1 hour. A saturated aqueous solution of ammonium chloride (50 ml) was added, and the mixture was extracted with ether. 10 The etheric layers were combined and dried over anhydrous magnesium sulfate. The dried product was filtered, concentrated, and subjected to silica gel column chromatography (silica gel, 50 g, eluent, benzene:ethyl acetate=10:1 \rightarrow 3:1) to give 329 mg 15 (yield 48%) of 3-n-butyl-4-butyldimethylsilyloxy-2-(1-hydroxybutyl)cyclopentanone.

The resulting product had the following spectral data.

IR (liquid film):

20 3450, 1735, 1250 cm^{-1} .

NMR (CDCl_3) δ :

0-0.2 (m, 6H), 0.7-1.1 (m, 6H), 0.89

(s, 9H), 1.1-3.0 (m, 15H), 3.6-4.5 (m, 2H).

(2) 137 mg (0.4 mmole) of 3-n-butyl-4-t-butyl-di- 25 methylsilyloxy-2-(1-hydroxybutyl)cyclopentanone obtained in (1) above was dissolved in a mixture of acetic acid, tetrahydrofuran and water in a ratio of 2:1:1, and the solution was refluxed at 70°C for 15 hours. Water and sodium bicarbonate were added to 30 render the reaction mixture basic, and it was extracted with ethyl acetate. The organic layers were combined, washed with a saturated aqueous solution of sodium chloride and dried over anhydrous magnesium sulfate. The dried product was filtered, concen- 35 trated, and subjected to silica gel column chromatography (silica gel, 5 g; eluent, hexane:ethyl acetate=5:1

- 37 -

- 1:1) to give 61 mg (yield 73%) of 4-n-butyl-5-(1-hydroxybutyl)-2-cyclopentenone. The spectral data of the resulting compound were as follows:-

IR (liquid film):

5 3480, 1698, 1585 cm^{-1} .

NMR (CDCl_3) δ :

0.7-1.1 (m, 6H), 1.1-3.0 (m, 13H),

3.7-4.1 (m, 1H), 6.22 (dd, 1H, J=5.8,

2.4Hz), 7.81 (dd, 1H, J=5.8m 2.8Hz).

10 Examples 16 to 19

(1) In the same way as in Example 15, (1) the 3-alkyl-4-t-butyldimethylsilyloxy-2-(1-hydroxyalkyl)-cyclopentanones shown in Table 1 were obtained.

Table 1

Ex-ample	Compound obtained	Yield (%)	IR (cm^{-1})	NMR (CDCl_3) δ
16	3-n-Butyl-4-t-butyl-4-dimethylsilyl-2-methylpropyl-cyclopentanone	64	3470 1735 1252	0-0.2 (m, 6H), 0.7-1.1 (m, 9H), 0.89 (s, 9H), 1.1-1.9 (m, 12H), 3.3-3.6 (m, 1H), 3.9-4.5 (m, 1H),
17	3-n-Butyl-4-t-butyl-4-dimethylsilyl-2,2-dimethyl-propyl)cyclopentanone	34	3490 1731 1250	0-0.2 (m, 6H), 0.7-1.1 (m, 3H), 0.89 (s, 18H), 1.1-2.7 (m, 11H), 3.1-3.5 (m, 1H), 4.3-4.7 (m, 1H)
18	3-n-Butyl-4-t-butyl-4-dimethylsilyl-2-oxy-2-(1-hydroxy-3-phenyl-2-propen-1-yl)cyclopentanone	45	3480 1723 1251 962	0-0.2 (m, 6H), 0.7-1.1 (m, 3H), 0.89 (s, 9H), 1.1-2.8 (m, 10H), 3.3-3.6 (m, 1H), 4.2-4.8 (m, 2H), 6.32 (dd, 1H, J=16.0, 6.8Hz), 6.60 (d, 1H, J=16.0Hz), 7.0-7.6 (m, 5H)
19	3-t-Butyl-4-t-butyl-4-dimethylsilyl-2-oxy-2-(1-hydroxybutyl)cyclopentanone	37	3490 1738 1249	0-0.2 (m, 6H), 0.7-1.1 (m, 3H), 0.90 (s, 18H), 1.1-2.7 (m, 9H), 3.3-3.6 (m, 1H), 4.0-4.5 (m, 1H)

- 39 -

(2) In the same way as in Example 15, (2) the 4-alkyl-5-(1-hydroxyalkyl)2-cyclopentenones indicated in Table 2 were obtained from the 3-alkyl-4-t-butyl-dimethylsilyloxy-2-(1-hydroxyalkyl)cyclopentenones
5 obtained in (1) above. The results are shown in Table 2.

Table 2

Ex-ample	Compound obtained	Yield (%)	IR ₁ (cm ⁻¹)	NMR (CDCl ₃) ^δ
16	4-n-Butyl-5-(1-hydroxy-2-methyl-propyl)-2-cyclopentanone	63	3510 1700 1580	0.7-1.1 (m, 9H), 1.1-3.1 (m, 10H), 3.4-3.8 (m, 1H), 6.20 (dd, 1H, J=5.8, 2.5Hz), 7.80 (dd, 1H, J=5.8, 2.8Hz)
17	4-n-Butyl-5-(1-hydroxy-2,2-dimethylpropyl)-2-cyclopentanone	41	3495 1696 1590	0.7-1.1 (m, 3H), 1.1-3.1 (m, 9H), 3.1-3.6 (m, 1H), 6.21 (dd, 1H, J=5.9, 2.6Hz), 7.80 (dd, 1H, J=5.9, 2.8Hz)
18	4-n-Butyl-5-(1-hydroxy-3-phenyl-2-propen-1-yl)-2-cyclopentanone	46	3490 1697 1595 960	0.7-1.1 (m, 3H), 1.1-3.4 (m, 7H), 3.3-3.6 (m, 1H), 4.3-4.7 (m, 1H), 6.0-6.8 (m, 3H), 7.0-8.0 (m, 6H)
19	4-t-Butyl-5-(1-hydroxybutyl)-2-cyclopentanone	40	3500 1701 1585	0.7-1.1 (m, 3H), 0.90 (s, 9H), 1.1-3.0 (m, 7H), 3.6-4.0 (m, 1H), 6.25 (dd, 1H, J=6.0, 2.6Hz), 7.75 (dd, 1H, J=6.0, 2.8Hz)

Example 20

5 Synthesis of 3-n-butyl-4-t-butyldimethylsilyloxy-2-(1-hydroxy-6-methoxycarbonylhexyl)cyclopentanone and 4-n-butyl-5-(1-hydroxy-6-methoxycarbonylhexyl)-2-cyclopentenone:-

(1) 6.2 ml (20 mmoles) of hexamethylphosphoric triamide was added to 1.88 g (14.4 mmoles) of 1-pentynyl copper, and the mixture was stirred for
10 30 minutes. Then, 50 ml of dry ether was added, and the mixture was cooled to -78°C . 9.6 ml (14.4 mmoles) of n-butyllithium (1.5M-hexane solution) was added, and the mixture was stirred for
15 15 minutes. A solution of 2.55 g (12 mmoles) of 4-t-butyldimethylsilyl-2-cyclopentenone in 30 ml of dry ether was added, and the mixture was stirred at -40°C for 15 minutes. A solution of 2.28 g (14.4 mmoles) of methyl 7-oxoheptanoate in 30 ml of dry ether was added, and the mixture was stirred at
20 -40°C for 1 hour. A saturated aqueous solution of ammonium chloride and ammonia and hexane were added to extract the reaction mixture. The organic layer was washed with a saturated aqueous solution of ammonium chloride and then with a saturated aqueous solution
25 of sodium chloride, and dried over anhydrous magnesium sulfate. The dried product was filtered, concentrated and subjected to silica gel column chromatography (silica gel 100 g; eluent, hexane:ethyl acetate=8:1 - 2:1) to give 0.55 g (yield 11%) of
30 3-n-butyl-4-butyldimethylsilyloxy-2-(1-hydroxy-6-methoxycarbonylhexyl)cyclopentanone. The resulting compound had the following spectral data.

IR (liquid film):

3500, 1740, 1250 cm^{-1} .

35 NMR (CDCl_3) δ :

0-0.2 (m, 6H), 0.7-1.1 (m, 3H), 0.89

- 42 -

(s, 9H), 1.1-2.8 (m, 21H), 3.70 (s, 3H),
3.7-4.4 (m, 2H).

(2) In the same way as in Example 15, (2) 4-n-butyl-5-(2-hydroxy-6-methoxycarbonylhexyl)-2-cyclopentenone was obtained in a yield of 73% from 3-n-butyl-4-t-butyldimethylsilyloxy-2-(1-hydroxy-6-methoxycarbonylhexyl)cyclopentanone obtained in (1) above. The resulting compound had the following spectral data.

10 IR (liquid film):

3480, 1737, 1698, 1585 cm^{-1} .

NMR (CDCl_3) δ :

0.7-1.1 (m, 3H), 1.1-3.0 (m, 20H), 3.70
(s, 3H), 3.7-4.1 (m, 1H), 6.21 (dd, 1H,
15 J=5.8, 2.4Hz), 7.81 (dd, 1H, J=5.8,
12.8Hz).

Example 21

Synthesis of 3-(2-propenyl)-4-t-butyldimethylsilyloxy-2-(1-hydroxybutyl)cyclopentanone and 4-(2-propenyl)-5-(1-hydroxybutyl)-2-cyclopentenone:-

(1) 266 mg (2.2 mmoles) of 2-bromopropene was dissolved in 5 ml of dry ether, and the solution was cooled to -95°C . Then, 3.0 ml (4.8 mmoles) of
20 t-butyllithium (as 1.60M pentane solution) was added, and the mixture was stirred at -78°C for 2 hours to form a 2-propenyllithium solution. 300 mg (2.05 mmoles) of cuprous iodide was taken into a reaction vessel. The reaction vessel was purged with argon,
25 and 30 ml of dry ether and 1.02 ml (4.1 mmoles) of tributyl phosphine were added. The mixture was stirred for 10 minutes. The mixture was added to the 2-propenyllithium solution prepared above, and the mixture was stirred for 5 minutes. A solution of 425
35 mg (2.0 mmoles) of 4-t-butyldimethylsilyloxy-2-cyclopentenone in 5 ml of dry ether was added. The mixture

- 43 -

was stirred at -78°C for 10 minutes and -40°C for 20 minutes. A solution of 148 mg (2.05 mmoles) of butanal in 5 ml of dry ether was added, and the mixture was stirred at -40°C for 1 hour. 50 ml of a saturated aqueous solution of ammonium chloride was added, and the mixture was extracted with ether. The etheric layers were combined, dried over anhydrous magnesium sulfate, filtered, concentrated and then subjected to silica gel column chromatography (silica gel 50 g; eluent, benzene:ethyl acetate=10:1 ~ 3:1 to give 287 mg (yield 44%) of 3-(2-propenyl)-4-t-butyl-dimethylsilyloxy-2-(1-hydroxybutyl)cyclopentanone. The resulting compound had the following spectral data.

IR (liquid film):

3480, 1737, 1640, 1251 cm^{-1} .

NMR (CDCl_3) δ :

0-0.2 (m, 6H), 0.7-1.1 (m, 3H), 0.90 (s, 3H), 1.1-3.3 (m, 12H), 3.6-4.5 (m, 2H).

(2) In the same way as in Example 15, (2) 4-(2-propenyl)-5-(1-hydroxybutyl)-2-cyclopentenone was obtained from 3-(2-propenyl)-4-t-butyl-dimethylsilyloxy-2-(1-hydroxybutyl)cyclopentanone obtained in (1) above. The resulting compound had the following spectral data.

IR (liquid film):

3490, 1696, 1637, 1586 cm^{-1} .

NMR (CDCl_3) δ :

0.7-1.1 (m, 3H), 1.1-2.8 (m, 9H), 3.0-3.5 (m, 1H), 3.7-4.1 (m, 1H), 6.22 (dd, 1H, $J=5.8, 2.3\text{Hz}$), 7.79 (dd, 1H, $J=5.8, 2.8\text{Hz}$).

Example 22

Synthesis of 4-t-butyl-dimethylsilyloxy-3-(3-t-butyl-dimethylsilyloxy-1-octenyl)-2-(1-hydroxy-6-methoxycarbonylhexyl)cyclopentanone and 4-(3-hydroxy-1-octenyl)-5-(1-hydroxy-6-methoxycarbonylhexyl)-2-

- 44 -

cyclopentenone:-

(1) 12 ml (24 mmoles) of t-butyllithium (2.0M pentane solution was cooled to -78°C , and 40 ml of dry ether was added. A solution of 4.42 g of 1-iodo-3-
5 t-butyldimethylsilyloxy-1-octene in 40 ml of dry ether, and the mixture was stirred at -78°C for 2 hours. A solution prepared by dissolving 1.566 g (12 mmoles) of 1-pentynyl copper (1) in 6.2 ml (29 mmoles) of hexamethylphosphoric triamide and
10 adding 20 ml of dry ether was added and the mixture was stirred at -78°C for 10 minutes. A solution of 2.12 g (10 mmoles) of 4-t-butyldimethylsilyloxy-2-cyclopentenone in 20 ml of dry ether was added. The mixture was stirred at -78°C for 5 minutes at
15 -40°C for 1 hour. The reaction mixture was poured into an acetic acid-sodium acetate buffer (pH 4), and hexane was added to perform extraction. The organic layer was washed with a saturated aqueous solution of sodium chloride, and dried over anhydrous magnesium
20 sulfate. The dried product was filtered, concentrated, and subjected to silica gel column chromatography (silica gel, 200 g; eluent, hexane:ethyl acetate=10:1 \rightarrow 2:1 to give 3.19 g (yield 52%) of 4-t-butyldimethylsilyloxy-3-(t-butyldimethylsilyloxy-1-octenyl)-2-(1-hydroxy-6-methoxycarbonylhexyl)-
25 cyclopentanone. The resulting compound had the following spectral data.

IR (liquid film):

3470, 1741, 1251 cm^{-1} .30 NMR (CDCl_3) δ :

0-0.2 (m, 12H), 0.7-1.1 (m, 3H), 0.90 (s, 18H), 1.1-3.0 (m, 23H), 3.69 (s, 3H), 3.6-4.6 (m, 3H), 5.4-5.8 (m, 2H).

(2) 1.23 g (2.0 mmoles) of 4-t-butyldimethylsilyloxy-3-(3-t-butyldimethylsilyloxy-1-octenyl)-2-(1-hydroxy-6-methoxycarbonylhexyl)cyclopentanone

35

- 45 -

obtained in (1) above was dissolved in 10 ml of a hydrogen fluoride-acetonitrile solution (composed of 47% hydrochloric acid and acetonitrile in a ratio of 1:20), and the solution was stirred for 30 minutes.

5 Sodium bicarbonate and water were added, and the mixture was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous solution of sodium chloride, and dried over anhydrous magnesium sulfate. The dried product was filtered,

10 concentrated, and subjected to silica gel column chromatography (silica gel 20 g; eluent, hexane:ethyl acetate=2:1 - 1:4) to give 577 mg of 4-hydroxy-3-(3-hydroxy-1-octenyl)-2-(1-hydroxy-6-methoxycarbonylhexyl)-cyclopentanone. This compound was further dissolved in

15 10 ml of a mixed solvent composed of acetic acid, tetrahydrofuran and water in a ratio of 2:1:1, and the solution was stirred at 70°C for 6 hours. An aqueous solution of sodium bicarbonate and ethyl acetate were added to perform extraction. The

20 organic layers were combined, washed with a saturated aqueous solution of sodium chloride, and dried over anhydrous magnesium chloride. The dried product was filtered, concentrated, and subjected to silica gel column chromatography (silica gel, 15 g; eluent,

25 hexane:ethyl acetate=5:1 - 1:2) to give 352 mg (yield 48%) of 4-(3-hydroxy-1-octenyl)-2-(1-hydroxy-6-methoxycarbonylhexyl)-2-cyclopentenone. The product had the following spectral data.

IR (liquid film):

30 3470, 1703, 1586 cm^{-1} .

NMR (CDCl_3) δ :

0.7-1.1 (m, 3H), 1.1-2.9 (m, 21H), 2.9-3.4 (m, 1H), 3.69 (s, 3H), 3.5-4.6 (m, 2H), 5.5-5.9 (m, 2H), 6.24 (dd, 1H, $J=5.8$, 2.1Hz),

35 7.63 (dd, 1H, $J=5.8$, 2.4Hz).

- 46 -

Example 23

43 mg (0.11 mmole) of 4-hydroxy-3-(3-hydroxy-1-octenyl)-2-(1-hydroxy-6-methoxycarbonylhexyl)cyclopentanone obtained as a reaction intermediate in
5 Example 22, (2) was dissolved in 1.5 ml of tetrahydrofuran, and 1.0 ml of 0.5N hydrochloric acid was added. The mixture was stirred at 40°C for 3 hours. An aqueous solution of sodium bicarbonate and ethyl acetate were added to perform extraction.
10 The organic layer was washed with a saturated aqueous solution of sodium chloride, and dried over anhydrous magnesium sulfate. The dried product was filtered, concentrated, and subjected to silica gel column chromatography (silica gel, 2 g; eluent, hexane:ethyl
15 acetate=5:1 → 1:2) to give 21 mg (yield 51%) of 4-(3-hydroxy-1-octenyl)-5-(1-hydroxy-6-methoxycarbonylhexyl)-2-cyclopentenone.

Example 24

370 mg (60 mmoles) of 4-t-butyldimethyl
20 silyloxy-3-(3-t-butyldimethylsilyloxy-1-octenyl)-2-(1-hydroxy-6-methoxycarbonylhexyl)cyclopentanone obtained in Example 22, (1) was dissolved in 10 ml of a mixed solvent consisting of acetic acid, tetrahydrofuran and water in a ratio of 2:1:1, and the solution
25 was stirred at 80°C for 7 hours. An aqueous solution of sodium bicarbonate and ethyl acetate were added to perform extraction. The organic layers were combined, washed with a saturated aqueous solution of sodium chloride, and dried over anhydrous
30 magnesium sulfate. The dried product was filtered, concentrated, and subjected to silica gel column chromatography (silica gel), 10 g; eluent, hexane:ethyl acetate=5:1 → 1:2) to give 218 mg (yield 62%) of
35 4-(3-hydroxy-1-octenyl)-5-(1-hydroxy-6-methoxycarbonylhexyl)cyclopentenone.

- 47 -

Examples 25 to 27

In the same way as in Example 15, the 3-alkenyl-4-t-butyldimethylsilyloxy-2-(1-hydroxyalkyl)cyclopentanones shown in Table 3 and the 4-alkenyl-5-(1-hydroxyalkyl)-2-cyclopentenones in Table 4 were obtained.

Table 3

Ex-ample	Compound obtained	Yield (%)	IR ₁ (cm ⁻¹)	NMR (CDCl ₃) ^δ
25	4-t-Butyldimethylsilyloxy-3-(3-t-butyl- dimethylsilyloxy-3-cyclo- pentyl-1-propenyl)-2-(1-hydroxy- 6-methoxycarbonyl-2-hexyn-1-yl)- cyclopentanone	45	3470 2230 1738 1252	0-0.2 (m, 12H), 0.89 (s, 18H), 1.2-3.0 (m, 20H), 3.70 (s, 3H), 3.5-4.5 (m, 2H), 4.6-5.0 (m, 1H), 5.5-5.8 (m, 2H)
26	4-(2-Tetrahydropranyloxy)-3-[3-(2-tetrahydropranyloxy)-1- octenyl]-2-(1-hydroxy-6-methoxy- carbonyl-2-hexyn-1-yl)cyclo- pentanone	15	3510 2230 1736	0.7-1.1 (m, 3H), 1.1-3.0 (m, 31H), 3.3-4.5 (m, 6H), 3.68 (s, 3H) 4.5-5.1 (m, 3H), 5.4-5.8 (m, 2H)
27	4-t-Butyldimethylsilyloxy-3-(3-t- butyldimethylsilyloxy-5-methyl- 1-nonenyl)-2-(1-hydroxy-6- methoxycarbonyl-5-hexen-1-yl)- cyclopentanone	42	3490 1735 1250	0-0.2 (m, 12H), 0.90 (s, 18H), 0.7-1.1 (m, 6H), 1.1-3.2 (m, 20H), 3.71 (s, 3H), 3.3-4.7 (m, 3H), 5.3-5.8 (m, 2H), 5.85 (brd, 1H, 16Hz), 7.03 (dt, 1H, J=16.0, 6.8Hz)

Table 4

Ex-ample	Compound obtained	Yield (%)	IR ₁ (cm ⁻¹)	NMR (CDCl ₃) ^δ
25	4-(3-t-Butyldimethylsilyloxy-3-cyclopentyl-1-propenyl)-5-(1-hydroxy-6-methoxycarbonyl-2-hexyn-1-yl)-2-cyclopentenone	26	3450 2230 1735 1708 1580 1247	0-0.2 (m, 6H), 0.89 (s, 9H), 1.2-2.7 (m, 18H), 3.1-3.5 (m, 1H), 3.68 (s, 3H), 3.7-4.1 (m, 1H), 4.6-5.0 (m, 1H), 5.5-5.8 (m, 2H), 6.22 (dd, 1H, J=5.8, 2.0Hz), 7.68 (dd, 1H, J=5.8, 3.2Hz)
26	4-[3-(2-tetrahydropyranyloxy)-1-octenyl]-5-(1-hydroxy-6-methoxycarbonyl-2-hexyn-1-yl)-2-cyclopentenone	36	3480 2230 1735 1706 1577	0.7-1.1 (m, 3H), 1.1-2.8 (m, 22H), 3.0-4.4 (m, 4H), 3.70 (s, 3H), 4.5-5.0 (m, 2H), 5.4-5.7 (m, 2H), 6.21 (dd, 1H, J=5.7, 2.0Hz), 7.67 (dd, 1H, J=5.7, 2.2Hz)
27	4-(3-t-Butyldimethylsilyloxy-5-methyl-1-nonenyl)-5-(1-hydroxy-6-methoxycarbonyl-5-hexen-1-yl)-2-cyclopentenone	12	3510 1734 1704 1250	0-0.2 (m, 6H), 0.89 (s, 9H), 0.7-1.1 (m, 6H), 1.1-2.7 (m, 17H), 2.9-3.4 (m, 1H), 3.72 (s, 3H), 3.55-4.6 (m, 2H), 5.5-5.9 (m, 2H), 5.88 (brd, 1H, J=16.0Hz), 6.25 (dd, 1H, J=5.8, 2.0Hz), 7.04 (dt, 1H, J=16.0, 7.2Hz), 7.63 (dd, 1H, J=5.8, 2.4Hz)

- 50 -

Example 28

Synthesis of 4-(3-hydroxy-5-methyl-1-nonenyl)-5-(1-hydroxy-6-methoxycarbonyl-5-hexen-1-yl)-2-cyclopentenone:-

5 In the same way as in Example 22, 6.7 mg (yield 43%) of 4-(3-hydroxy-5-methyl-1-nonenyl)-5-(1-hydroxy-6-methoxycarbonyl-5-hexen-1-yl)-2-cyclopentenone was obtained from 20 mg (0.039 mmole) of 4-(3-t-butyltrimethylsilyloxy-5-methyl-1-nonenyl)-
10 5-(1-hydroxy-6-methoxycarbonyl-5-hexen-1-yl)-2-cyclopentenone obtained in Example 27. The resulting compound had the following spectral data.

IR (liquid film):

3520, 1702, 1580 cm^{-1} .

15 NMR (CDCl_3) δ :

0.7-1.05 (m, 6H), 1.05-2.0 (m, 13H), 2.0-2.7 (m, 5H), 2.9-3.5 (m, 1H), 3.74 (s, 3H), 3.55-4.6 (m, 2H), 5.5-5.9 (m, 2H), 5.88 (brd, 1H, $J=16.0\text{Hz}$), 6.26 (dd, 1H, $J=5.8$, 2.0Hz), 7.05 (dt, 1H, $J=16.0$, 7.2Hz), 7.64
20 7.64 (dd, 1H, $J=5.8$, 2.4Hz).

Example 29

Synthesis of 4-(3-hydroxy-3-cyclopentyl-1-propenyl)-5-(1-hydroxy-6-methoxycarbonyl-2-hexyn-1-yl)-2-cyclopentenone:-

25 63 mg (0.133 mmole) of 4-(3-t-butyltrimethylsilyloxy-3-cyclopentyl-1-propenyl)-5-(1-hydroxy-6-methoxycarbonyl-2-hexyn-1-yl)-2-cyclopentenone obtained in Example 25 was dissolved in 2 ml of a mixed
30 solvent composed of acetic acid, tetrahydrofuran and water in a ratio of 3:2:2, and the solution was stirred for 20 hours. An aqueous solution of sodium bicarbonate and ethyl acetate were added to the reaction mixture to perform extraction. The organic
35 layers were combined, washed with a saturated aqueous solution of sodium chloride, and dried over anhydrous

- 51 -

sodium sulfate. The dried product was filtered, concentrated and subjected to silica gel column chromatography (silica gel 5 g; eluent, hexane:ethyl acetate=5:1 - 1:1) to give 38 mg (yield 79%) of 4-(3-hydroxy-3-cyclopentyl-1-propenyl)-5-(1-hydroxy-6-methoxycarbonyl-2-hexynyl)-2-cyclopentenone. The resulting compound had the following spectral data.

IR (liquid film):

3450, 2240, 1700, 1583 cm^{-1} .

10 NMR (CDCl_3) δ :

0.9-2.8 (m, 18H), 3.68 (s, 3H), 3.5-4.3 (m, 3H), 4.6-5.0 (m, 1H), 5.6-6.0 (m, 2H), 6.21 (dd, 1H, $J=6.0, 2.0\text{Hz}$), 7.71 (dd, 1H, $J=6.0, 2.4\text{Hz}$).

15 Example 30

Synthesis of 4-(3-hydroxy-1-octenyl)-5-(1-hydroxy-6-carboxyhexyl)-2-cyclopentenone:-

Acetone (0.6 ml) and 6 ml of phosphate buffer (pH 8) were added to 50 mg (0.135 mmole) of 4-(3-hydroxy-1-octenyl)-5-(1-hydroxy-6-methoxycarbonylhexyl)-2-cyclopentenone obtained in Example 24, and then 0.06 ml of an aqueous solution of esterase (from pig liver) was added. The mixture was stirred for 25 70 hours. Hydrochloric acid was added to the reaction mixture to adjust its pH to 4, and then the mixture was saturated with ammonium sulfate. It was filtered, and then extracted with ethyl acetate. The organic layer was washed with an aqueous solution of sodium chloride, and dried over anhydrous magnesium sulfate. The dried product was filtered, concentrated and subjected to silica gel column chromatography (silica gel 2 g; eluent, hexane:ethyl acetate=2:1 - 1:4) to give 31 mg (yield 65%) of 4-(3-hydroxy-1-octenyl)-5-(1-hydroxy-6-carboxylhexyl)-2-cyclopentene. 35 The resulting compound had the following spectral data

- 52 -

IR (liquid film):

3400, 1702, 1585 cm^{-1} .NMR (CDCl_3) δ :

0.7-1.1 (m, 3H), 1.1-2.9 (m, 22H), 2.9-3.4
5 (m, 1H), 3.5-4.6 (m, 2H), 5.4-5.9 (m, 2H),
6.23 (dd, 1H, $J=5.8$, 2.1Hz), 7.63 (dd, 1H,
 $J=5.8$, 2.4Hz)

Example 31

Synthesis of 4-(3-t-butyldimethylsilyloxy-
10 1-octenyl)-5-(6-methoxycarbonylhexylidene)-
2-cyclopentenone:-

480 mg (1.0 mmole) of 4-(3-t-butyldimethyl-
silyloxy-1-octenyl)-5-(1-hydroxy-6-methoxycarbonyl-
hexyl)-2-cyclopentenone was dissolved in 5 ml of
15 dichloromethane, followed by addition of 730 mg
(6 mmoles) of dimethylaminopyridine and 230 micro-
liters (3.0 mmoles) of methanesulfonyl chloride in
this order. The mixture was stirred at 40°C for 2
hours. An aqueous solution of sodium bicarbonate was
20 added, and the mixture was stirred with dichloro-
methane. The organic layers were combined, washed
with a saturated aqueous solution of sodium chloride
and dried over anhydrous sodium sulfate. The dried
product was filtered, concentrated and subjected to
25 silica gel column chromatography (silica gel 20 g;
eluent, hexane:ethyl acetate=10:1) to give 114 mg
(yield 25%) of 4-(3-t-butyldimethylsilyloxy-1-octenyl)-
5-(6-methoxycarbonylhexylidene)-2-cyclopentenone.
The resulting compound had the following spectral
30 data.

TLC: $R_f=0.54$ (hexane:ethyl acetate=3:1)NMR (CDCl_3) δ :

0-0.2 (m, 6H), 0.86 (s, 9H), 0.7-1.0
(m, 3H), 1.1-2.0 (m, 14H), 2.0-2.7
35 (m, 4H), 3.67 (s, 3H), 3.7-4.3 (m, 2H),
5.3-5.8 (m, 2H), 6.36 (dd, 1H, $J=6.0$,

- 53 -

1.8Hz), 6.65 (brt, 1H, J=8.0Hz), 7.45
(dd, 1H, J=6.0, 2.8Hz).

Example 32

- Synthesis of 4-butyl-5-(6-methoxycarbonyl-
hexylidene)-2-cyclopentenone:-
- 1.7 g (5.74 mmoles) of 4-butyl-5-(1-hydroxy-
6-methoxycarbonyl)-2-cyclopentenone was dissolved in
20 ml of dichloromethane, and 3.86 g (31.6 mmoles) of
4-dimethylaminopyridine was added. With ice cooling
and stirring, 900 microliters (11.6 mmoles) of methane-
sulfonyl chloride was added. The mixture was stirred
at 0°C for 10 minutes and then at room temperature
for 20 hours. Water and hydrochloric acid was added
to adjust the pH of the reaction mixture to 1, and it
was extracted with methylene chloride. The organic
layer was washed with a saturated aqueous solution of
sodium bicarbonate, and dried over anhydrous sodium
sulfate. The dried product was filtered, concen-
trated and subjected to silica gel column chromato-
graphy (silica gel, 20 g; eluent, hexane:ethyl
acetate=8:1 → 4:1) to give 111 mg (yield 7%) of
4-butyl-(Z)-5-(6-methoxycarbonylhexylidene)-2-cyclo-
pentenone and 1.363 g (yield 86%) of 4-butyl-(Z)-5-(6-
methoxycarbonylhexylidene)-2-cyclopentenone.
- Spectral data of 4-butyl-(Z)-5-(6-methoxy-
carbonylhexylidene)-2-cyclopentenone:-
TLC: Rf=0.53 (hexane:ethyl acetate=2:1)
IR(liquid film):
1739, 1694, 1642, 1583 cm⁻¹.
- NMR (CDCl₃)δ:
0.7-1.1 (m, 3H), 1.1-2.0 (m, 12H), 2.0-2.55
(m, 2H), 2.55-3.2 (m, 2H), 3.1-3.5 (m, 1H),
3.69 (s, 3H), 6.07 (brt, 1H, J=7.5Hz), 6.31
(dd, 1H, J=6.0, 2.0Hz), 7.50 (dd, 1H, J=6.0,
3.0Hz).

- 54 -

Spectral data of 4-butyl-(E)-5-(6-methoxycarbonylhexylidene)-2-cyclopentenone:-

TLC: Rf=0.42 (hexane:ethyl acetate=2:1)

IR (liquid film):

5 1739, 1703, 1656, 1580 cm^{-1} .

NMR (CDCl_3) δ :

0.89 (brt, 3H), 1.0-2.0 (m, 12H), 2.0-2.6 (m, 4H), 3.3-3.8 (m, 1H), 3.67 (s, 3H)
6.35 (dd, 1H, J=6.0, 2.0Hz), 6.56 (brt, 1H),
10 7.59 (dd, 1H, J=6.0, 3.0Hz).

Example 33

Synthesis of 4-(1-octenyl)-5-(1-hydroxy-6-methoxycarbonylhexylidene)-2-cyclopentenone:-

15 In the same way as in Example 15, 4-(1-octenyl)-5-(1-hydroxy-6-methoxycarbonylhexylidene)-2-cyclopentenone was obtained in an amount of 1.32 g (yield 38%). The product had the following spectral data.

TLC: Rf=0.13 (hexane:ethyl acetate=4:1)

NMR (CDCl_3) δ :

20 7.86 (brt, 3H, J=4.5Hz), 1.0-1.8 (m, 16H), 1.85-2.5 (m, 5H), 3.1-3.4 (m, 1H), 3.64 (s, 3H), 3.4-4.0 (m, 1H), 4.0-4.2 (m, 1H), 7.55 (dd, 1H, J=6.0, 2.4Hz).

Example 34

25 Synthesis of 4-(1-octenyl)-5-(6-methoxycarbonylhexylidene)-2-cyclopentenone:-

300 mg (0.86 mmole) of 4-(1-octenyl)-5-(1-hydroxy-6-methoxycarbonylhexyl)-2-cyclopentene obtained in Example 33 was dissolved in 5 ml of anhydrous dichloromethane, and 640 mg (5.2 mmoles) of dimethylaminopyridine and 200 microliters (2.6 mmoles) of methanesulfonyl chloride were added in this order. The mixture was stirred at room temperature for 4 hours, and 50 ml of ether was added. The organic layer was
35 washed with an acetic buffer (pH 4) and then with water and further with an aqueous solution of sodium

- 55 -

chloride, and dried. The solvent was evaporated, and the residue was purified by thin-layer chromatography (solvent, hexane:ether=3:1, R_f =0.4, R_f of the starting enone being 0.25) to give 135 mg (yield 48%) of a
5 mixture of isomers of 4-(1-octenyl)-5-(6-methoxycarbonyl-hexylidene)-2-cyclopentenone of Example 5.

Example 35

Measurement of the action of inhibiting proliferation of L1210 leukemia cell:-

10 L1210 leukemia cells were added rto an RPMI medium containing 10% FCS (fetal calf serum), and the concentration of the cells was adjusted to 1×10^5 cells/ml. Each of the test compounds shown in Table 5 was dissolved in 99.5% ethanol. Prior to use, the
15 final concentration of the ethanol solution was adjusted to less than 0.1%, and it was added to the culture medium. The culture medium was then maintained at 37°C in a stationary condition for 4 days. After the cultivation, the number of surviv-
20 ing cells was measured by dyeing with trypan blue. As a control, 0.1% ethanol was used. A dose-reaction curve was plotted from the ratios of proliferation against the control, and LC_{50} was determined.

The results are shown in Table 5.

Table 5

Test compound	IC ₅₀ (μg/ml)
7(E)-7,8-Dehydroprostaglandin E ₁	0.7
7(E)-7,8-Dehydroprostaglandin A ₁	0.4
4-Butyl-5-(6-methoxycarbonylhexylidene)-2-cyclopentenone	0.3
4-(3-Hydroxy-3-cyclopentyl-1-propenyl)-5-(6-methoxycarbonyl-2-hexynylidene)-2-cyclopentenone	0.3
4-(3-Hydroxy-5-methyl-1-nonenyl)-5-(6-methoxycarbonyl-2-hexynylidene)-2-cyclopentenone [7(E)-7,8-dehydro-17(S),20-dimethylprostaglandin A ₁ methyl ester]	0.2
4-(3-Hydroxy-3-cyclopentyl-1-propenyl)-5-(1-hydroxy-6-methoxycarbonyl-2-hexyn-1-yl)-2-cyclopentenone	0.2
12-Epi-(7E)-7,8-dehydroprostaglandin A ₁ methyl ester	0.2
4-(1-Octenyl)-5-(6-methoxycarbonyl-hexynylidene)-2-cyclopentenone	0.3

- 57 -

Example 36

Measurement of the antitumor effect on Ehrlich ascites carcinoma:-

1 x 10⁵ Ehrlich ascites carcinoma cells were
5 intraperitoneally administered to ICR mice. After the
lapse of 24 hours, 30 mg/kg/day of 7(E)-7,8-dehydroprosta-
glandin A₁ methyl ester and 20 mg/kg/day of its
12-epimer were each intraperitoneally administered to
the mice for 5 days. The periods of survival of
10 these animals were examined.

When 7(E)-7,8-dehydroprostaglandin A₁ methyl
ester was administered, the average number of days of
survival was 31±1.9 days. The increase of life
span (ILS %) increased by 65.8% over the control, and
15 the ratio of survival for more than 60 days was 2/6.

In the case of administering its 120 epimer,
the average number of days of survival was 33.0±
9.8 days. The increase of life span (ILS %) in-
creased by 76.5% over the control, and the ratio of
20 survival for more than 60 days was 1/6.

Example 37

Measurement of cyto protection:-

Stomach epithelial cells taken from a rabbit
fetus and aortic smooth muscle cells from a rat were
25 used as normal cells, and Ll210 leukemia cells were
used as cancer cells. The cytotoxicity of each of
the compounds shown in Table 6 on the normal cells
was examined.

Specifically, the stomach epithelial cells
30 were cultivated in DME containing 20% FCS, and the
aortic smooth muscle cells were cultivated in MEM
containing 10% CS. Each of the test compounds was
dissolved in ethanol and added to the culture broth
in an amount of 0.1%. The mixture was filtered
35 through a millipore filter.

The cytotoxicities (LD₅₀) of each of the

- 58 -

test compounds on the stomach epithelial cells and the aortic smooth muscle cells were determined. The results are shown in Table 6. Furthermore, from the LFD₅₀ values, the safety coefficient (LD₅₀ on the
5 normal cells/LD₅₀ on the L1210 cells) was calculated, and the results are also shown in Table 6.

Table 6

Test compound	LD ₅₀ (µg/ml)			Safety coefficient
	Stomach epithelial cells	Aortic smooth muscle cells	L1210 leukemia cells	
7(E)-7,8-dehydroprostaglandin A ₁	2 - 5	2 - 5	0.2 - 0.3	6 - 25
12-Epimer of 7(E)-7,8-dehydroprostaglandin A ₁	1 - 2	1 - 2	0.2 - 0.4	2.5 - 10
Comparison (Mycromycin C)	0.01 - 0.02	0.01	0.1	0.01 - 0.1

- 60 -

It is seen from Table 6 that the compounds of this invention have low cytotoxicity on normal cells.

Example 38

Measurement of acute toxicity:-

- 5 The acute toxicity of one typical compound of this invention was measured by a customary method using four weeks old LR strain male mice (SPF).

The results are shown in Table 7.

Table 7

Compound	Administration route	LD ₅₀ (mg/kg)
7(E)-7,8-dehydroprostaglandin A ₁	i.v.	140

- 10 Table 7 shows that the compound of this invention has low acute toxicity.

Example 39

Production of soft capsules:-

- 15 One milligrams of (7E)-7,8-dehydro PGA₁ obtained in Example 2 was dissolved in 60 g of fractionated coconut oil and soft capsules were produced using a soft gelatin capsule making machine. Each of the capsules containing 1 g of (7E)-7,8-dehydro PGA₁.

20 Example 40

Production of tablets:-

Tablets were produced each of which contained the following ingredients.

- | | | |
|----|--|--------|
| 25 | 4-(3-hydroxy-3-cyclopentyl-1-propenyl)-5-(1-hydroxy-6-methoxycarbonyl-2-hexyn-1-yl)-2-cyclopentenone | 10 g |
| | Lactonse | 250 mg |

- 61 -

Potato starch	70 mg
Polyvinyl pyrrolidone	10 mg
Magnesium stearate	5 mg

The aforesaid cyclopentenone compound obtained
5 in Example 29, lactose and potato starch were mixed,
and the mixture was evenly wetted with a 20% ethanol
solution of polyvinyl pyrrolidone. The wetted mi-
xture was passed through a sieve. The resulting
granules were mixed with magnesium stearate, and
10 compression-molded into tablets.

Example 41

Preparation of an injectable solution:-

(7E)-7,8-dehydro PGA₁ obtained in Example 2
as an active ingredient was dissolved in an amount of
15 60 µg in 5 ml of ethanol, and the solution was steri-
lized by passing it through a bacteria-holding
filter. 0.1 ml of the solution was filled in each of
1-ml ampoules and the ampoules were sealed up. The
contents of the ampoules are diluted, for example,
20 with Tris-HCl buffer to 1 ml for injection.

Example 42

Production of a powde:-

A powder was prepared in accordance with the
following recipe.

25	4-(1-Octenyl)-5-(6-methoxy- carbonylhexylidene)-2-cyclo- pentanone	10 µg
	Lactose	100 mg
	Corn starch	100 mg
	Hydroxypropyl cellulose	10 mg

The above cyclopentenone compound obtained as
30 a typical active ingredient, lactose and corn starch
were mixed, and an aqueous solution of hydroxypropyl
cellulose was added. The mixture was dried to form a
powder.

- 62 -

CLAIMS

1. A 4,5-disubstituted-2-cyclopentenone selected from 5-alkylidene-4-substituted-2-cyclopentenones represented by formula (I)




wherein W represents an aliphatic hydrocarbon group having 1 to 12 carbon atoms which may have a substituent, and Y represents an aliphatic hydrocarbon group having 1 to 12 carbon atoms which may have a substituent, and 5-(1-hydroxy-aliphatic hydrocarbon)-4-substituted-2-cyclopentenones represented by formula (II)



wherein W' and Y' are as defined for W and Y respectively in formula (I).


2. A 4,5-disubstituted-2-cyclopentenone according to claim 1 wherein the compounds of formula (I) are 5-alkylidene-4-substituted-2-cyclopentenones represented by formula (I)-1



- 63 -

wherein R^1 represents a hydrogen atom or an aliphatic hydrocarbon group having 1 to 10 carbon atoms which may have a substituent, R^2 represents a hydrogen atom or an aliphatic hydrocarbon group having 1 to 9 carbon atoms which may have a substituent, R^3 represents a hydrogen atom, a hydroxyl group or a protected hydroxyl group, and the symbol $\begin{smallmatrix} \cdot & \cdot & \cdot & \cdot \\ \text{---} & & \text{---} & \\ \cdot & \cdot & \cdot & \cdot \end{smallmatrix}$ represents a single, double or triple bond.

3. A 4,5-disubstituted-2-cyclopentenone according to claim 1 wherein the compounds of formula (II) are 5-(1-hydroxy-aliphatic hydrocarbons)-4-substituted-2-cyclopentenones represented by formula (II)-1



... (II)-1


wherein R^1 , R^2 , R^3 and the symbol $\begin{smallmatrix} \cdot & \cdot & \cdot & \cdot \\ \text{---} & & \text{---} & \\ \cdot & \cdot & \cdot & \cdot \end{smallmatrix}$ are as defined in claim 2.

4. A 4,5-disubstituted-2-cyclopentenone according to any one of claims 1 to 3 wherein the substituent on the aliphatic hydrocarbon group represented by W, W', Y or Y' is a group of the formula $-\text{COOR}^4$ in which R^4 represents a hydrogen atom, an alkyl group having 1 to 10 carbon atoms or one equivalent of a cation; a group of the formula $-\text{OR}^5$ in which R^5 represents a hydrogen atom, an alkyl group having 1 to 6 carbon atoms which may be substituted by a halogen atom, a carboacyl group having 1 to 7 carbon atoms or a phenyl group, the phenyl group being optionally substituted by a halogen atom, an alkyl group having 1 to 4 carbon atoms or an alkoxy group having 1 to 4 carbon atoms; a phenyl group which may be substituted by a halogen atom, an alkyl group

- 64 -


having 1 to 4 carbon atoms or an alkoxy group having 1 to 4 carbon atoms; or a cycloalkyl group having 3 to 8 carbon atoms which may be substituted by a halogen atom, an alkyl group having 1 to 4 carbon atoms or an alkoxy group having 1 to 4 carbon atoms.

5. A 4,5-disubstituted-2-cyclopentenone according to claim 1 or 2 which is a 5-alkylidene-4-substituted-2-cyclopentenone represented by formula (I)-2



wherein R^2 , R^3 , R^4 and the symbol $\text{---}\cdot\cdot\cdot\cdot$ are as hereinbefore defined, and the symbol $\text{---}\cdot\cdot\cdot\cdot$ represents a single or double bond.

6. A 4,5-disubstituted-2-cyclopentenone according to claim 1 or 3 which is a 5-(1-hydroxy-hydrocarbon)-4-substituted-2-cyclopentenone represented by formula (II)-2



wherein R^2 , R^3 , R^4 , and the symbols $\text{---}\cdot\cdot\cdot\cdot$ and $\text{---}\cdot\cdot\cdot\cdot$ are as hereinbefore defined.

7. A process for producing a 5-alkylidene-4-substituted-2-cyclopentenone of formula (I) as claimed in claim 1 which comprises subjecting

- 65 -


a 5-alkylidene-3-hydroxy-4-substituted-cyclopentanone represented by formula (III)



wherein W' and Y' are as defined for W and Y respectively,

to a dehydration reaction, and as required, subjecting the reaction product to a deprotecting, hydrolyzing or salt-forming reaction.

8. A process for producing a 5-(1-hydroxy-aliphatic hydrocarbon)-4-substituted-2-cyclopentenone of formula (II) as claimed in claim 1 which comprises subjecting a 5-(1-hydroxy-aliphatic hydrocarbon)-3-hydroxy-4-substituted cyclopentanone represented by formula (IV)



wherein W'' and Y'' are the same as defined for W and Y respectively,

to a dehydration reaction, and as required, subjecting the reaction product to a deprotecting, hydrolyzing or salt-forming reaction.

- 66 -

9. A pharmaceutical composition comprising, as active ingredient, a 5-membered cyclic compound selected from 5-alkylidene-4-substituted-2-cyclopentenones represented by formula (I) as defined in claim 1, 5-(alpha-hydroxy-aliphatic hydrocarbon)-4-substituted-2-cyclopentenones represented by formula (II) as defined in claim 1 and 5-alkylidene-3-hydroxy-4-substituted cyclopentanones represented by formula (III) as defined in claim 7 and a pharmaceutically acceptable carrier.

10. A pharmaceutical composition according to claim 9 for use in the treatment of tumors.

11. A medicament in unit dosage form comprising a pharmaceutical composition as claimed in claim 9 or 10.

THIS PAGE BLANK (USPTO)